



Ad: Study [discontinuous analysis](#), an enhanced calculus in which every function is both differentiable and integrable.

Ad: [World-best general purpose programming language](#). You won't like Python anymore.

Ad: [Donate for science](#).

# Algebraic General Topology. Volume 1 addons

Victor Porton

*Email address:* [porton@narod.ru](mailto:porton@narod.ru)

*URL:* <http://www.mathematics21.org>

2000 *Mathematics Subject Classification.* 54J05, 54A05, 54D99, 54E05, 54E15,  
54E17, 54E99

*Key words and phrases.* algebraic general topology, quasi-uniform spaces,  
generalizations of proximity spaces, generalizations of nearness spaces,  
generalizations of uniform spaces

ABSTRACT. This file contains future addons for the free e-book “Algebraic  
General Topology. Volume 1”, which are yet not enough ripe to be included  
into the book.

# Contents

Chapter 1. About this document	6
Chapter 2. Applications of algebraic general topology	7
1. “Hybrid” objects	7
2. A way to construct directed topological spaces	7
3. Some inequalities	9
4. Continuity	10
5. A way to construct directed topological spaces	13
6. Integral curves	13
Chapter 3. Covers	18
Chapter 4. More on generalized limit	19
1. Hausdorff funcoids	19
2. Restoring functions from limit	19
Chapter 5. Extending Galois connections between funcoids and reloids	21
Chapter 6. Boolean funcoids	23
1. One-element boolean lattice	23
2. Two-element boolean lattice	23
3. Finite boolean lattices	24
4. About infinite case	24
Chapter 7. Interior funcoids	26
Chapter 8. Filterization of pointfree funcoids	28
Chapter 9. Systems of sides	29
1. More on Galois connections	29
2. Definition	30
3. Concrete examples of sides	31
4. Product	33
5. Negative results	34
6. Dagger systems of sides	34
Chapter 10. Backward Funcoids	35
Chapter 11. Quasi-atoms	36
Chapter 12. Cauchy Filters on Reloids	37
1. Preface	37
2. Low spaces	37
3. Almost sub-join-semilattices	38
4. Cauchy spaces	38
5. Relationships with symmetric reloids	39
6. Lattices of low spaces	40

7. Up-complete low spaces	44
8. More on Cauchy filters	45
9. Maximal Cauchy filters	46
10. Cauchy continuous functions	47
11. Cauchy-complete reloids	47
12. Totally bounded	47
13. Totally bounded funcoids	48
14. On principal low spaces	48
15. Rest	48
Chapter 13. Funcoidal groups	50
1. On “Each regular paratopological group is completely regular” article	51
Chapter 14. Micronization	56
Chapter 15. More on connectedness	57
1. For topological spaces	57
Chapter 16. Relationships are pointfree funcoids	62
Chapter 17. Manifolds and surfaces	63
1. Sides of a surface	63
2. Special points	63
Bibliography	66

## CHAPTER 1

### **About this document**

This file contains future addons for the free e-book “Algebraic General Topology. Volume 1”, which are yet not enough ripe to be included into the book.

Theorem (including propositions, conjectures, etc.) numbers in this document start from the last theorem number in the book plus one. Theorems references inside this document are hyperlinked, but references to theorems in the book are not hyperlinked (because PDF viewer Okular 0.20.2 does not support Backward button after clicking a cross-document reference, and thus I want to avoid clicking such links).

## Applications of algebraic general topology

### 1. “Hybrid” objects

Algebraic general topology allows to construct “hybrid” objects of “continuous” (as topological spaces) and discrete (as graphs).

Consider for example  $D \sqcup T$  where  $D$  is a digraph and  $T$  is a topological space.

The  $n$ -th power  $(D \sqcup T)^n$  yields an expression with  $2^n$  terms. So treating  $D \sqcup T$  as one object (what becomes possible using algebraic general topology) rather than the join of two objects may have an exponential benefit for simplicity of formulas.

### 2. A way to construct directed topological spaces

**2.1. Some notation.** I use  $\mathcal{E}$  and  $\iota$  notations from `volume-2.pdf`. FiXme: Reorder document fragments to describe it before use.

I remind that  $f|_X = f \circ \text{id}_X$  for binary relations, funcoids, and reloid.

$$f \parallel_X = f \circ (\mathcal{E}^X)^{-1}.$$

$$f \square X = \text{id}_X \circ f \circ \text{id}_X^{-1}.$$

As proved in `volume-2.pdf`, the following are bijections and moreover isomorphisms (for  $R$  being either funcoids or reloids or binary relations):

$$1^\circ. \left\{ \frac{(f|_X, f \parallel_X)}{f \in R} \right\};$$

$$2^\circ. \left\{ \frac{(f \square X, \iota_X f)}{f \in R} \right\}.$$

As easily follows from these isomorphisms and theorem 1182:

PROPOSITION 2111. For funcoids, reloids, and binary relations:

$$1^\circ. f \in C(\mu, \nu) \Rightarrow f \parallel_A \in C(\iota_A \mu, \nu);$$

$$2^\circ. f \in C'(\mu, \nu) \Rightarrow f \parallel_A \in C'(\iota_A \mu, \nu);$$

$$3^\circ. f \in C''(\mu, \nu) \Rightarrow f \parallel_A \in C''(\iota_A \mu, \nu).$$

**2.2. Directed line and directed intervals.** Let  $\mathfrak{A}$  be a poset. We will denote  $\overline{\mathfrak{A}} = \mathfrak{A} \cup \{-\infty, +\infty\}$  the poset with two added elements  $-\infty$  and  $+\infty$ , such that  $+\infty$  is strictly greater than every element of  $\mathfrak{A}$  and  $-\infty$  is strictly less.

FiXme: Generalize from  $\mathbb{R}$  to a wider class of posets.

DEFINITION 2112. For an element  $a$  of a poset  $\mathfrak{A}$

$$1^\circ. J_{\geq}(a) = \left\{ \frac{x \in \mathfrak{A}}{x \geq a} \right\};$$

$$2^\circ. J_{>}(a) = \left\{ \frac{x \in \mathfrak{A}}{x > a} \right\};$$

$$3^\circ. J_{\leq}(a) = \left\{ \frac{x \in \mathfrak{A}}{x \leq a} \right\};$$

$$4^\circ. J_{<}(a) = \left\{ \frac{x \in \mathfrak{A}}{x < a} \right\};$$

$$5^\circ. J_{\neq}(a) = \left\{ \frac{x \in \mathfrak{A}}{x \neq a} \right\}.$$

DEFINITION 2113. Let  $a$  be an element of a poset  $\mathfrak{A}$ .

$$1^\circ. \Delta(a) = \prod^{\mathcal{F}} \left\{ \frac{[x; y]}{x, y \in \mathfrak{A}, x < a \wedge y > a} \right\};$$

$$2^\circ. \Delta_{\geq}(a) = \prod^{\mathcal{F}} \left\{ \frac{[a; y]}{y \in \mathfrak{A}, y > a} \right\};$$

$$\begin{aligned}
3^\circ. \Delta_{>}(a) &= \prod^{\mathcal{F}} \left\{ \frac{]a;y[}{y \in \mathfrak{A}, x < a \wedge y > a} \right\}; \\
4^\circ. \Delta_{\leq}(a) &= \prod^{\mathcal{F}} \left\{ \frac{]x;a[}{x \in \mathfrak{A}, x < a} \right\}; \\
5^\circ. \Delta_{<}(a) &= \prod^{\mathcal{F}} \left\{ \frac{]x;a[}{x \in \mathfrak{A}, x < a} \right\}; \\
6^\circ. \Delta_{\neq}(a) &= \Delta(a) \setminus \{a\}.
\end{aligned}$$

OBVIOUS 2114.

$$\begin{aligned}
1^\circ. \Delta_{\geq}(a) &= \Delta(a) \sqcap^{\mathcal{F}} @J_{\geq}(a); \\
2^\circ. \Delta_{>}(a) &= \Delta(a) \sqcap^{\mathcal{F}} @J_{>}(a); \\
3^\circ. \Delta_{\leq}(a) &= \Delta(a) \sqcap^{\mathcal{F}} @J_{\leq}(a); \\
4^\circ. \Delta_{<}(a) &= \Delta(a) \sqcap^{\mathcal{F}} @J_{<}(a); \\
5^\circ. \Delta_{\neq}(a) &= \Delta(a) \sqcap^{\mathcal{F}} @J_{\neq}(a).
\end{aligned}$$

DEFINITION 2115. Given a partial order  $\mathfrak{A}$  and  $x \in \mathfrak{A}$ , the following defines complete functors:

$$\begin{aligned}
1^\circ. \langle |\mathfrak{A}| \rangle^* \{x\} &= \Delta(x); \\
2^\circ. \langle |\mathfrak{A}|_{\geq} \rangle^* \{x\} &= \Delta_{\geq}(x); \\
3^\circ. \langle |\mathfrak{A}|_{>} \rangle^* \{x\} &= \Delta_{>}(x); \\
4^\circ. \langle |\mathfrak{A}|_{\leq} \rangle^* \{x\} &= \Delta_{\leq}(x); \\
5^\circ. \langle |\mathfrak{A}|_{<} \rangle^* \{x\} &= \Delta_{<}(x); \\
6^\circ. \langle |\mathfrak{A}|_{\neq} \rangle^* \{x\} &= \Delta_{\neq}(x).
\end{aligned}$$

PROPOSITION 2116. The complete functor corresponding to the order topology<sup>1</sup> is equal to  $|\mathfrak{A}|$ .

PROOF. Because every open set is a finite union of open intervals, the complete functor  $f$  corresponding to the order topology is described by the formula:  $\langle f \rangle^* \{x\} = \prod^{\mathcal{F}} \left\{ \frac{]a;b[}{a, b \in \mathfrak{A}, a < x \wedge b > x} \right\} = \Delta(x) = \langle |\mathfrak{A}| \rangle^* \{x\}$ . Thus  $f = |\mathfrak{A}|$ .  $\square$

EXERCISE 2117. Show that  $|\mathfrak{A}|_{\geq}$  (in general) is not the same as “right order topology”<sup>2</sup>.

PROPOSITION 2118.

$$\begin{aligned}
1^\circ. \langle |\mathfrak{A}|_{\geq}^{-1} \rangle^* @X &= @ \left\{ \frac{a \in \mathfrak{A}}{\forall y \in \mathfrak{A}: (y > a \Rightarrow X \cap ]a;y[ \neq \emptyset)} \right\}; \\
2^\circ. \langle |\mathfrak{A}|_{>}^{-1} \rangle^* @X &= @ \left\{ \frac{a \in \mathfrak{A}}{\forall y \in \mathfrak{A}: (y > a \Rightarrow X \cap ]a;y[ \neq \emptyset)} \right\}; \\
3^\circ. \langle |\mathfrak{A}|_{\leq}^{-1} \rangle^* @X &= @ \left\{ \frac{a \in \mathfrak{A}}{\forall x \in \mathfrak{A}: (x < a \Rightarrow X \cap ]x;a[ \neq \emptyset)} \right\}; \\
4^\circ. \langle |\mathfrak{A}|_{<}^{-1} \rangle^* @X &= @ \left\{ \frac{a \in \mathfrak{A}}{\forall x \in \mathfrak{A}: (x < a \Rightarrow X \cap ]x;a[ \neq \emptyset)} \right\}.
\end{aligned}$$

PROOF.  $a \in \langle |\mathfrak{A}|_{\geq}^{-1} \rangle^* @X \Leftrightarrow @\{a\} \neq \langle |\mathfrak{A}|_{\geq}^{-1} \rangle^* @X \Leftrightarrow \langle |\mathfrak{A}|_{\geq} \rangle^* @\{a\} \neq @X \Leftrightarrow \Delta_{\geq}(a) \neq @X \Leftrightarrow \forall y \in \mathfrak{A}: (y > a \Rightarrow X \cap ]a;y[ \neq \emptyset)$ .

$a \in \langle |\mathfrak{A}|_{>}^{-1} \rangle^* @X \Leftrightarrow @\{a\} \neq \langle |\mathfrak{A}|_{>}^{-1} \rangle^* @X \Leftrightarrow \langle |\mathfrak{A}|_{>} \rangle^* @\{a\} \neq @X \Leftrightarrow \Delta_{>}(a) \neq @X \Leftrightarrow \forall y \in \mathfrak{A}: (y > a \Rightarrow X \cap ]a;y[ \neq \emptyset)$ .

The rest follows from duality.  $\square$

REMARK 2119. On trivial ultrafilters these obviously agree:

$$\begin{aligned}
1^\circ. \langle |\mathbb{R}|_{\geq} \rangle^* \{x\} &= \langle |\mathbb{R}| \cap \geq \rangle^* \{x\}; \\
2^\circ. \langle |\mathbb{R}|_{>} \rangle^* \{x\} &= \langle |\mathbb{R}| \cap > \rangle^* \{x\}; \\
3^\circ. \langle |\mathbb{R}|_{\leq} \rangle^* \{x\} &= \langle |\mathbb{R}| \cap \leq \rangle^* \{x\}; \\
4^\circ. \langle |\mathbb{R}|_{<} \rangle^* \{x\} &= \langle |\mathbb{R}| \cap < \rangle^* \{x\}.
\end{aligned}$$

<sup>1</sup>See Wikipedia for a definition of “Order topology”.

<sup>2</sup>See Wikipedia



COROLLARY 2120.

- 1°.  $|\mathbb{R}|_{\geq} = \text{Compl}(|\mathbb{R}| \cap \geq)$ ;
- 2°.  $|\mathbb{R}|_{>} = \text{Compl}(|\mathbb{R}| \cap >)$ ;
- 3°.  $|\mathbb{R}|_{\leq} = \text{Compl}(|\mathbb{R}| \cap \leq)$ ;
- 4°.  $|\mathbb{R}|_{<} = \text{Compl}(|\mathbb{R}| \cap <)$ .

OBVIOUS 2121. **FiXme:** also what is the values of  $\setminus$  operation

- 1°.  $|\mathbb{R}|_{\geq} = |\mathbb{R}|_{>} \sqcup 1$ ;
- 2°.  $|\mathbb{R}|_{\leq} = |\mathbb{R}|_{<} \sqcup 1$ .

### 3. Some inequalities

**FiXme:** Define the ultrafilter “at the left” and “at the right” of a real number. Also define “convergent ultrafilter”.

Denote  $\Delta_{+\infty} = \prod_{x \in \mathbb{R}} x; +\infty[$  and  $\Delta_{-\infty} = \prod_{x \in \mathbb{R}} ] - \infty; x[$ .

The following proposition calculates  $\langle \geq \rangle x$  and  $\langle > \rangle x$  for all kinds of ultrafilters on  $\mathbb{R}$ :

PROPOSITION 2122.

- 1°.  $\langle \geq \rangle \{\alpha\} = [\alpha; +\infty[$  and  $\langle > \rangle \{\alpha\} = ]\alpha; +\infty[$ .
- 2°.  $\langle \geq \rangle x = \langle > \rangle x = ]\alpha; +\infty[$  for ultrafilter  $x$  at the right of a number  $\alpha$ .
- 3°.  $\langle \geq \rangle x = \langle > \rangle x = \Delta_{<}(\alpha) \sqcup [\alpha; +\infty[ = \Delta_{\leq}(\alpha) \sqcup ]\alpha; +\infty[$  for ultrafilter  $x$  at the left of a number  $\alpha$ .
- 4°.  $\langle \geq \rangle x = \langle > \rangle x = \Delta_{+\infty}$  for ultrafilter  $x$  at positive infinity.
- 5°.  $\langle \geq \rangle x = \langle > \rangle x = \mathbb{R}$  for ultrafilter  $x$  at negative infinity.

PROOF.

- 1°. Obvious.
- 2°.

$$\begin{aligned} \langle \geq \rangle x &= \prod_{X \in \text{up } x}^{\mathcal{F}} \langle \geq \rangle (X \cap \alpha; +\infty[) = \prod_{X \in \text{up } x}^{\mathcal{F}} ]\alpha; +\infty[ = ]\alpha; +\infty[; \\ \langle > \rangle x &= \prod_{X \in \text{up } x}^{\mathcal{F}} \langle > \rangle (X \cap \alpha; +\infty[) = \prod_{X \in \text{up } x}^{\mathcal{F}} ]\alpha; +\infty[ = ]\alpha; +\infty[. \end{aligned}$$

- 3°.  $\Delta_{<}(\alpha) \sqcup [\alpha; +\infty[ = \Delta_{\leq}(\alpha) \sqcup ]\alpha; +\infty[$  is obvious.

$$\langle > \rangle x = \prod_{X \in \text{up } x}^{\mathcal{F}} \langle > \rangle X \supseteq \prod_{X \in \text{up } x}^{\mathcal{F}} (\Delta_{<}(\alpha) \sqcup ]\alpha; +\infty[) = \Delta_{<}(\alpha) \sqcup ]\alpha; +\infty[$$

but  $\langle \geq \rangle x \subseteq \Delta_{<}(\alpha) \sqcup [\alpha; +\infty[$  is obvious. It remains to take into account that  $\langle > \rangle x \subseteq \langle \geq \rangle x$ .

$$\begin{aligned} 4°. \quad \langle \geq \rangle x &= \prod_{X \in \text{up } x}^{\mathcal{F}} \langle \geq \rangle X = \prod_{X \in \text{up } x, \inf X \in X}^{\mathcal{F}} \langle \geq \rangle (X \cap \alpha; +\infty[) = \\ &= \prod_{X \in \text{up } x}^{\mathcal{F}} [\inf X; +\infty[ = \prod_{x > \alpha}^{\mathcal{F}} [x; +\infty[ = \Delta_{+\infty}; \quad \langle > \rangle x = \prod_{X \in \text{up } x}^{\mathcal{F}} \langle > \rangle X = \\ &= \prod_{X \in \text{up } x, \inf X \in X}^{\mathcal{F}} \langle > \rangle (X \cap \alpha; +\infty[) = \prod_{X \in \text{up } x}^{\mathcal{F}} \inf X; +\infty[ = \prod_{x > \alpha}^{\mathcal{F}} [x; +\infty[ = \Delta_{+\infty}. \end{aligned}$$

- 5°.  $\langle \geq \rangle x \supseteq \langle > \rangle x = \prod_{X \in \text{up } x}^{\mathcal{F}} \langle > \rangle X$  but  $\langle > \rangle X = ] - \infty; +\infty[$  for  $X \in \text{up } x$  because  $X$  has arbitrarily small elements.

□

LEMMA 2123.  $\langle |\mathbb{R}| \rangle x \subseteq \langle > \rangle x = \langle \geq \rangle x$  for every nontrivial ultrafilter  $x$ .

PROOF.  $\langle > \rangle x = \langle \geq \rangle x$  follows from the previous proposition.

$$\langle |\mathbb{R}| \rangle x = \prod_{X \in \text{up } x} \langle |\mathbb{R}| \rangle X = \prod_{X \in \text{up } x} \bigsqcup_{y \in X} \Delta(y).$$

Consider cases:

$x$  is an ultrafilter at the right of some number  $\alpha$ .

$$\langle |\mathbb{R}| \rangle x = \prod_{X \in \text{up } x} \bigsqcup_{y \in X \cap ]\alpha; +\infty[} \Delta(y) \sqsubseteq ]\alpha; +\infty[ = \langle \geq \rangle x \quad \text{because} \\ \bigsqcup_{y \in X \cap ]\alpha; +\infty[} \Delta(y) \sqsubseteq ]\alpha; +\infty[.$$

$x$  is an ultrafilter at the left of some number  $\alpha$ .

$$\langle |\mathbb{R}| \rangle x \sqsubseteq \Delta(\alpha) \text{ is obvious. But } \langle \geq \rangle x \supseteq \Delta(\alpha).$$

$x$  is an ultrafilter at positive infinity.

$$\langle |\mathbb{R}| \rangle x \sqsubseteq \Delta_{+\infty} \text{ is obvious. But } \langle \geq \rangle x = \Delta_{+\infty}.$$

$x$  is an ultrafilter at negative infinity.

$$\text{Because } \langle \geq \rangle x = \mathbb{R}.$$

□

COROLLARY 2124.  $\langle |\mathbb{R}| \cap \geq \rangle x = \langle |\mathbb{R}| \rangle x$  for every nontrivial ultrafilter  $x$ .

$$\text{PROOF. } \langle |\mathbb{R}| \cap \geq \rangle x = \langle |\mathbb{R}| \rangle \cap \langle \geq \rangle x = \langle |\mathbb{R}| \rangle x.$$

□

So  $\langle |\mathbb{R}| \cap \geq \rangle$  and  $\langle |\mathbb{R}| \rangle$  agree on all ultrafilters except trivial ones.

PROPOSITION 2125.  $|\mathbb{R}|_{>} \cap > = |\mathbb{R}|_{>} \cap \geq = |\mathbb{R}|_{>}$ .

PROOF.  $|\mathbb{R}|_{>} \sqsubseteq >$  because  $\langle |\mathbb{R}|_{>} \rangle^* x \sqsubseteq \langle > \rangle^* x$  and  $|\mathbb{R}|_{>}$  is a complete functor.

□

LEMMA 2126.  $\langle |\mathbb{R}|_{>} \rangle x \sqsubset \langle |\mathbb{R}|_{\geq} \rangle x$  for a nontrivial ultrafilter  $x$ .

PROOF. It enough to prove  $\langle |\mathbb{R}|_{>} \rangle x \neq \langle |\mathbb{R}|_{\geq} \rangle x$ .

Take  $x$  be an ultrafilter with limit point 0 on  $\text{im } z$  where  $z$  is the sequence  $n \mapsto \frac{1}{n}$ .

$$\langle |\mathbb{R}|_{>} \rangle x \sqsubseteq \langle |\mathbb{R}|_{>} \rangle^* \text{im } z = \bigsqcup_{n \in \text{im } z} \Delta_{>} \left( \frac{1}{n} \right) \sqsubseteq \bigsqcup_{n \in \text{im } z} \left] \frac{1}{n}; \frac{1}{n-1} - \frac{1}{n} \right[ \asymp \text{im } z.$$

Thus  $\langle |\mathbb{R}|_{>} \rangle x \asymp \text{im } z$ . But  $\langle |\mathbb{R}|_{\geq} \rangle x \sqsubseteq \langle = \rangle x \not\asymp \text{im } z$ .

□

COROLLARY 2127.  $|\mathbb{R}|_{>} \sqsubset |\mathbb{R}|_{\geq}$ .

PROPOSITION 2128.  $|\mathbb{R}|_{>} \sqsubset |\mathbb{R}|_{\geq} \cap >$ .

PROOF. It's enough to prove  $|\mathbb{R}|_{>} \neq |\mathbb{R}|_{\geq} \cap >$ .

Really,  $\langle |\mathbb{R}|_{\geq} \cap > \rangle x = \langle |\mathbb{R}|_{\geq} \rangle x \neq \langle |\mathbb{R}|_{>} \rangle x$  (lemma).

□

PROPOSITION 2129.

- 1°.  $|\mathbb{R}|_{\geq} \circ |\mathbb{R}|_{\geq} = |\mathbb{R}|_{\geq}$ ;
- 2°.  $|\mathbb{R}|_{>} \circ |\mathbb{R}|_{>} = |\mathbb{R}|_{>}$ ;
- 3°.  $|\mathbb{R}|_{\geq} \circ |\mathbb{R}|_{>} = |\mathbb{R}|_{>}$ ;
- 4°.  $|\mathbb{R}|_{>} \circ |\mathbb{R}|_{\geq} = |\mathbb{R}|_{>}$ .

PROOF. ??

□

CONJECTURE 2130.

- 1°.  $(|\mathbb{R}| \cap \geq) \circ (|\mathbb{R}| \cap \geq) = |\mathbb{R}| \cap \geq$ .
- 2°.  $(|\mathbb{R}| \cap >) \circ (|\mathbb{R}| \cap >) = |\mathbb{R}| \cap >$ .

#### 4. Continuity

I will say that a property holds on a filter  $\mathcal{A}$  iff there is  $A \in \text{up } \mathcal{A}$  on which the property holds.

FixMe:  $f \in C(A, B) \wedge f \in C(\iota_A |\mathbb{R}|_{\geq}, \iota_B |\mathbb{R}|_{\geq}) \Leftrightarrow (f, f) \in C((A, \iota_A |\mathbb{R}|_{\geq}), (B, \iota_B |\mathbb{R}|_{\geq}))$

LEMMA 2131. Let function  $f : A \rightarrow B$  where  $A, B \in \mathcal{P}\mathbb{R}$  and  $\iota_A|\mathbb{R}|$  is connected.

- 1°.  $f$  is monotone and  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|)$  iff  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|) \cap C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq})$  iff  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|) \cap C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq})$  iff  $f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq}) \cap C(\iota_A|\mathbb{R}|_{\leq}, \iota_B|\mathbb{R}|_{\leq})$ .
- 2°.  $f$  is strictly monotone and  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|)$  iff  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|) \cap C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{>})$  iff  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{>}) \cap C(\iota_A|\mathbb{R}|_{<}, \iota_B|\mathbb{R}|_{<})$ .

FiXme: Generalize for arbitrary posets. FiXme: Generalize for  $f$  being a funcoid.  
FiXme: Can add more conditions with  $<$ .

PROOF. Because  $f$  is continuous, we have  $\langle f \circ \iota_A|\mathbb{R}| \rangle^* \{x\} \sqsubseteq \langle \iota_B|\mathbb{R}| \circ f \rangle^* \{x\}$  that is  $\langle f \rangle^*(A \cap \Delta(x)) \sqsubseteq B \cap \Delta(f(x))$  for every  $x \in A$ .

If  $f$  is monotone, we have  $\langle f \rangle^* \Delta_{\geq}(x) \sqsubseteq [f(x); \infty[$ . Thus  $\langle f \rangle^*(A \cap \Delta_{\geq}(x)) \sqsubseteq B \cap \Delta_{\geq}(f(x))$ , that is  $\langle f \circ \iota_A|\mathbb{R}|_{\geq} \rangle^* \{x\} \sqsubseteq \langle \iota_B|\mathbb{R}|_{\geq} \circ f \rangle^* \{x\}$ , thus  $f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq})$ .

If  $f$  is strictly monotone, we have  $\langle f \rangle^* \Delta_{>}(x) \sqsubseteq ]f(x); \infty[$ . Thus  $\langle f \rangle^* \Delta_{>}(x) \sqsubseteq \Delta_{>}(f(x))$ , that is  $\langle f \circ \iota_A|\mathbb{R}|_{>} \rangle^* \{x\} \sqsubseteq \langle \iota_B|\mathbb{R}|_{>} \circ f \rangle^* \{x\}$ , thus  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{>})$ .

Let now  $f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq})$ .

Take any  $a \in A$  and let  $c = \sup \left\{ \frac{b \in B}{b \geq a, \forall x \in [a; b[ : f(x) \geq f(a)} \right\}$  (makes sense because  $A$  is connected). It's enough to prove that  $c$  is the right endpoint (finite or infinite) of  $A$ .

Indeed by continuity  $f(a) \leq f(c)$  and if  $c$  is not already the right endpoint of  $A$ , then there is  $b' > c$  such that  $\forall x \in [c; b'[ : f(x) \geq f(c)$  (makes sense because  $A$  is connected). So we have  $\forall x \in [a; b'[ : f(x) \geq f(c)$  what contradicts to the above.

So  $f$  is monotone on the entire  $A$ .

$f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq}) \Rightarrow f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq})$  is obvious. Reversely  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq}) \Leftrightarrow f \circ \iota_A|\mathbb{R}|_{>} \sqsubseteq \iota_B|\mathbb{R}|_{\geq} \circ f \Leftrightarrow \forall x \in A : \langle f \rangle \langle \iota_A|\mathbb{R}|_{>} \rangle^* \{x\} \sqsubseteq \langle \iota_B|\mathbb{R}|_{\geq} \rangle^* \langle f \rangle^* \{x\} \Leftrightarrow \forall x \in A : \langle f \rangle (A \cap \Delta_{>}(x)) \sqsubseteq B \cap \Delta_{\geq} f(x) \Leftrightarrow \forall x \in A : \langle f \rangle (A \cap \Delta_{>}(x)) \sqcup \{f(x)\} \sqsubseteq B \cap \Delta_{\geq} f(x) \Leftrightarrow \forall x \in A : \langle f \rangle ((A \cap \Delta_{>}(x)) \sqcup \{x\}) \sqsubseteq B \cap \Delta_{\geq} f(x) \Leftrightarrow \forall x \in A : \langle f \rangle (A \cap \Delta_{\geq}(x)) \sqsubseteq B \cap \Delta_{\geq} f(x) \Leftrightarrow \forall x \in A : \langle f \rangle \langle \iota_A|\mathbb{R}|_{\geq} \rangle^* \{x\} \sqsubseteq \langle \iota_B|\mathbb{R}|_{\geq} \rangle^* \langle f \rangle^* \{x\} \Leftrightarrow f \circ \iota_A|\mathbb{R}|_{\geq} \sqsubseteq \iota_B|\mathbb{R}|_{\geq} \circ f \Leftrightarrow f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq})$ .

Let  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq})$ . Then  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq})$  and thus it is monotone. We need to prove that  $f$  is strictly monotone. Suppose the contrary. Then there is a nonempty interval  $[p; q] \subseteq A$  such that  $f$  is constant on this interval. But this is impossible because  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq})$ .

Prove that  $f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq}) \cap C(\iota_A|\mathbb{R}|_{\leq}, \iota_B|\mathbb{R}|_{\leq})$  implies  $f \in C(A, B)$ . Really, it implies  $\langle f \rangle (A \cap \Delta_{\leq}(x)) \sqsubseteq B \cap \Delta_{\leq}(f(x))$  and  $\langle f \rangle (A \cap \Delta_{\geq}(x)) \sqsubseteq B \cap \Delta_{\geq}(f(x))$  thus  $\langle f \rangle (A \cap \Delta(x)) = \langle f \rangle (A \cap (\Delta_{\leq}(x) \sqcup \{x\} \sqcup \Delta_{\geq}(x))) \sqsubseteq B \cap (\Delta_{\leq} f(x) \sqcup \{f(x)\} \sqcup \Delta_{\geq} f(x)) = B \cap \Delta(f(x))$ .

Prove that  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{>}) \cap C(\iota_A|\mathbb{R}|_{<}, \iota_B|\mathbb{R}|_{<})$  implies  $f \in C(A, B)$ . Really, it implies  $\langle f \rangle (A \cap \Delta_{<}(x)) \sqsubseteq B \cap \Delta_{<}(f(x))$  and  $\langle f \rangle (A \cap \Delta_{>}(x)) \sqsubseteq B \cap \Delta_{>}(f(x))$  thus  $\langle f \rangle (A \cap \Delta(x)) = \langle f \rangle (A \cap (\Delta_{<}(x) \sqcup \{x\} \sqcup \Delta_{>}(x))) \sqsubseteq B \cap (\Delta_{<} f(x) \sqcup \{f(x)\} \sqcup \Delta_{>} f(x)) = B \cap \Delta(f(x))$ .  $\square$

THEOREM 2132. FiXme: Counterexample: <https://math.stackexchange.com/a/3702872/4876> Let function  $f : A \rightarrow B$  where  $A, B \in \mathcal{P}\mathbb{R}$ .

- 1°.  $f$  is locally monotone and  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|)$  iff  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|) \cap C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq})$  iff  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|) \cap C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{\geq})$  iff  $f \in C(\iota_A|\mathbb{R}|_{\geq}, \iota_B|\mathbb{R}|_{\geq}) \cap C(\iota_A|\mathbb{R}|_{\leq}, \iota_B|\mathbb{R}|_{\leq})$ .

- 2°.  $f$  is locally strictly monotone and  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|)$  iff  $f \in C(\iota_A|\mathbb{R}|, \iota_B|\mathbb{R}|) \cap C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{>})$  iff  $f \in C(\iota_A|\mathbb{R}|_{>}, \iota_B|\mathbb{R}|_{>}) \cap C(\iota_A|\mathbb{R}|_{<}, \iota_B|\mathbb{R}|_{<})$ .

PROOF. By the lemma it is (strictly) monotone on each connected component.

FiXme: It is not enough if for example  $A = \mathbb{Q}$ .  $\square$

See also related math.SE questions:

- 1°. <http://math.stackexchange.com/q/1473668/4876>  
 2°. <http://math.stackexchange.com/a/1872906/4876>  
 3°. <http://math.stackexchange.com/q/1875975/4876>

**4.1. Directed topological spaces.** Directed topological spaces are defined at <http://ncatlab.org/nlab/show/directed+topological+space>

DEFINITION 2133. A *directed topological space* (or *d-space* for short) is a pair  $(X, d)$  of a topological space  $X$  and a set  $d \subseteq C([0; 1], X)$  (called *directed paths* or *d-paths*) of paths in  $X$  such that

- 1°. (constant paths) every constant map  $[0; 1] \rightarrow X$  is directed;
- 2°. (reparameterization)  $d$  is closed under composition with increasing continuous maps  $[0; 1] \rightarrow [0; 1]$ ;
- 3°. (concatenation)  $d$  is closed under path-concatenation: if the d-paths  $a, b$  are consecutive in  $X$  ( $a(1) = b(0)$ ), then their ordinary concatenation  $a + b$  is also a d-path

$$(a + b)(t) = a(2t), \text{ if } 0 \leq t \leq \frac{1}{2},$$

$$(a + b)(t) = b(2t - 1), \text{ if } \frac{1}{2} \leq t \leq 1.$$

I propose a new way to construct a directed topological space. My way is more geometric/topological as it does not involve dealing with particular paths.

DEFINITION 2134. Let  $T$  be the complete endofunctor corresponding to a topological space and  $\nu \sqsubseteq T$  be its “subfunctor”. The d-space  $(\text{dir})(T, \nu)$  induced by the pair  $(T, \nu)$  consists of  $T$  and paths  $f \in C([0; 1], T) \cap C([0; 1]_{\geq}, \nu)$  such that  $f(0) = f(1)$ .

PROPOSITION 2135. It is really a d-space.

PROOF. Every d-path is continuous.

Constant path are d-paths because  $\nu$  is reflexive.

Every reparameterization is a d-path because they are  $C([0; 1]_{\geq}, \nu)$  and we can apply the theorem about composition of continuous functions.

Every concatenation is a d-path. Denote  $f_0 = \lambda t \in [0; \frac{1}{2}] : a(2t)$  and  $f_1 = \lambda t \in [\frac{1}{2}; 1] : b(2t - 1)$ . Obviously  $f_0, f_1 \in C([0; 1], \mu) \cap C([0; 1]_{\geq}, \nu)$ . Then we conclude that  $a + b = f_0 \sqcup f_1$  is in  $f_0, f_1 \in C([0; 1], \mu) \cap C([0; 1]_{\geq}, \nu)$  using the fact that the operation  $\circ$  is distributive over  $\sqcup$ .  $\square$

Below we show that not every d-space is induced by a pair of an endofunctor and its subfunctor. But are d-spaces not represented this way good anything except counterexamples?

Let now we have a d-space  $(X, d)$ . Define functor  $\nu$  corresponding to the d-space by the formula  $\nu = \bigsqcup_{a \in d} (a \circ |\mathbb{R}|_{\geq} \circ a^{-1})$ .

EXAMPLE 2136. The two directed topological spaces, constructed from a fixed topological space and two different reflexive functors, are the same.

PROOF. Consider the indiscrete topology  $T$  on  $\mathbb{R}$  and the functors  $1^{\text{FCD}(\mathbb{R}, \mathbb{R})}$  and  $1^{\text{FCD}(\mathbb{R}, \mathbb{R})} \sqcup (\{0\} \times^{\text{FCD}} \Delta_{\geq})$ . The only d-paths in both these settings are constant functions.  $\square$

EXAMPLE 2137. A d-space is not determined by the induced functor.

PROOF. The following a d-space induces the same functor as the d-space of all paths on the plane.

Consider a plane  $\mathbb{R}^2$  with the usual topology. Let d-paths be paths lying inside a polygonal chain (in the plane).  $\square$

CONJECTURE 2138. A d-path  $a$  is determined by the functors (where  $x$  spans  $[0; 1]$ )

$$(\lambda t \in \mathbb{R} : a(x + t))|_{\Delta(0)}.$$

## 5. A way to construct directed topological spaces

I propose a new way to construct a directed topological space. My way is more geometric/topological as it does not involve dealing with particular paths.

CONJECTURE 2139. Every directed topological space can be constructed in the below described way.

Consider topological space  $T$  and its subfunctor  $F$  (that is  $F$  is a functor which is less than  $T$  in the order of functors). Note that in our consideration  $F$  is an endofunctor (its source and destination are the same).

Then a directed path from point  $A$  to point  $B$  is defined as a continuous function  $f$  from  $[0; 1]$  to  $F$  such that  $f(0) = A$  and  $f(1) = B$ . **FixMe: Specify whether the interval  $[0; 1]$  is treated as a proximity, pretopology, or preclosure.**

Because  $F$  is less than  $T$ , we have that every directed path is a path.

CONJECTURE 2140. The two directed topological spaces, constructed from a fixed topological space and two different functors, are different.

For a counter-example of (which of the two?) the conjecture consider functor  $T \sqcap (\mathbb{Q} \times^{\text{FCD}} \mathbb{Q})$  where  $T$  is the usual topology on real line. We need to consider stability of existence and uniqueness of a path under transformations of our functor and under transformations of the vector field. Can this be a step to solve Navier-Stokes existence and smoothness problems?

## 6. Integral curves

We will consider paths in a normed vector space  $V$ .

DEFINITION 2141. Let  $D$  be a connected subset of  $\mathbb{R}$ . A *path* is a function  $D \rightarrow V$ .

Let  $d$  be a vector field in a normed vector space  $V$ .

DEFINITION 2142. *Integral curve* of a vector field  $d$  is a differentiable function  $f : D \rightarrow V$  such that  $f'(t) = d(f(t))$  for every  $t \in D$ .

DEFINITION 2143. The definition of *right side integral curve* is the above definition with right derivative of  $f$  instead of derivative  $f'$ . *Left side integral curve* is defined similarly.

**6.1. Path reparameterization.**  $C^1$  is a function which has continuous derivative on every point of the domain.

By  $D^1$  I will denote a  $C^1$  function whose derivative is either nonzero at every point or is zero everywhere.

DEFINITION 2144. A *reparameterization* of a  $C^1$  path is a bijective  $C^1$  function  $\phi : D \rightarrow D$  such that  $\phi'(t) > 0$ . A curve  $f_2$  is called a reparametrized curve  $f_1$  if there is a reparameterization  $\phi$  such that  $f_2 = f_1 \circ \phi$ .

It is well known that this defines an equivalence relation of functions.

PROPOSITION 2145. Reparameterization of  $D^1$  function is  $D^1$ .

PROOF. If the function has zero derivative, it is obvious.

Let  $f_1$  has everywhere nonzero derivative. Then  $f_2'(t) = f_1'(\phi(t))\phi'(t)$  what is trivially nonzero.  $\square$

DEFINITION 2146. Vectors  $p$  and  $q$  have the *same direction* ( $p \uparrow\uparrow q$ ) iff there exists a strictly positive real  $c$  such that  $p = cq$ .

OBVIOUS 2147. Being same direction is an equivalence relation.

OBVIOUS 2148. The only vector with the same direction as the zero vector is zero vector.

THEOREM 2149. A  $D^1$  function  $y$  is some reparameterization of a  $D^1$  integral curve  $x$  of a continuous vector field  $d$  iff  $y'(t) \uparrow\uparrow d(y(t))$  that is vectors  $y'(t)$  and  $d(y(t))$  have the same direction (for every  $t$ ).

PROOF. If  $y$  is a reparameterization of  $x$ , then  $y(t) = x(\phi(t))$ . Thus  $y'(t) = x'(\phi(t))\phi'(t) = d(x(\phi(t)))\phi'(t) = d(y(t))\phi'(t)$ . So  $y'(t) \uparrow\uparrow d(y(t))$  because  $\phi'(t) > 0$ .

Let now  $x'(t) \uparrow\uparrow d(x(t))$  that is that is there is a strictly positive function  $c(t)$  such that  $x'(t) = c(t)d(x(t))$ .

If  $x'(t)$  is zero everywhere, then  $d(x(t)) = 0$  and thus  $x'(t) = d(x(t))$  that is  $x$  is an Integral curve. Note that  $y$  is a reparameterization of itself.

We can assume that  $x'(t) \neq 0$  everywhere. Then  $F(x(t)) \neq 0$ . We have that  $c(t) = \frac{\|x'(t)\|}{\|d(x(t))\|}$  is a continuous function. **FiXme: Does it work for non-normed spaces?**

Let  $y(u(t)) = x(t)$ , where

$$u(t) = \int_0^t c(s)ds,$$

which is defined and finite because  $c$  is continuous and monotone (thus having inverse defined on its image) because  $c$  is positive.

Then

$$\begin{aligned} y'(u(t))u'(t) &= x'(t) \\ &= c(t)d(x(t)), \text{ so} \\ y'(u(t))c(t) &= c(t)d(y(u(t))) \\ y'(u(t)) &= d(y(u(t))) \end{aligned}$$

and letting  $s = u(t)$  we have  $y'(s) = d(y(s))$  for a reparameterization  $y$  of  $x$ .  $\square$

**6.2. Vector space with additional coordinate.** Consider the normed vector space with additional coordinate  $t$ .

Our vector field  $d(t)$  induces vector field  $\hat{d}(t, v) = (1, d(v))$  in this space. Also  $\hat{f}(t) = (t, f(t))$ .

PROPOSITION 2150. Let  $f$  be a  $D^1$  function.  $f$  is an integral curve of  $d$  iff  $\hat{f}$  is a reparametrized integral curve of  $\hat{d}$ .

PROOF. First note that  $\hat{f}$  always has a nonzero derivative.  $\hat{f}'(t) \uparrow\uparrow \hat{d}(\hat{f}(t)) \Leftrightarrow (1, f'(t)) \uparrow\uparrow (1, d(f(t))) \Leftrightarrow f'(t) = d(f(t)). \quad \square$

Thus we have reduced (for  $D^1$  paths) being an integral curve to being a reparametrized integral curve. We will also describe being a reparametrized integral curve topologically (through funcoids).

**6.3. Topological description of  $C^1$  curves.** Explicitly construct this funcoid as follows:

$R(d, \phi) = \left\{ \frac{v \in V}{\widehat{vd} < \phi, v \neq 0} \right\}$  for  $d \neq 0$  and  $R(0, \phi) = \{0\}$ , where  $\widehat{ab}$  is the angle between the vectors  $a$  and  $b$ , for a direction  $d$  and an angle  $\phi$ .

DEFINITION 2151.  $W(d) = \prod_{\phi \in \mathbb{R}, \phi > 0}^{\text{RLD}} \left\{ \frac{R(d, \phi)}{\phi \in \mathbb{R}, \phi > 0} \right\} \sqcap \prod_{r > 0}^{\text{RLD}} B_r(0)$ . **FixMe:** This is defined for infinite dimensional case. **FixMe:** Consider also FCD instead of RLD.

PROPOSITION 2152. For finite dimensional case  $\mathbb{R}^n$  we have  $W(d) = \prod_{\phi \in \mathbb{R}, \phi > 0}^{\text{RLD}} \left\{ \frac{R(d, \phi)}{\phi \in \mathbb{R}, \phi > 0} \right\} \sqcap \Delta^{(\text{RLD})n}$  where

$$\Delta^{(\text{RLD})n} = \underbrace{\Delta \times^{\text{RLD}} \dots \times^{\text{RLD}} \Delta}_{n \text{ times}}.$$

PROOF. ??  $\square$

Finally our funcoids are the complete funcoids  $Q_+$  and  $Q_-$  described by the formulas

$$\langle Q_+ \rangle^* @ \{p\} = \langle p+ \rangle W(d(p)) \quad \text{and} \quad \langle Q_- \rangle^* @ \{p\} = \langle p+ \rangle W(-d(p)).$$

Here  $\Delta$  is taken from the ‘‘counter-examples’’ section.

In other words,

$$Q_+ = \bigsqcup_{p \in \mathbb{R}} (\@ \{p\} \times^{\text{FCD}} \langle p+ \rangle W(d(p))); \quad Q_- = \bigsqcup_{p \in \mathbb{R}} (\@ \{p\} \times^{\text{FCD}} \langle p+ \rangle W(-d(p))).$$

That is  $\langle Q_+ \rangle^* @ \{p\}$  and  $\langle Q_- \rangle^* @ \{p\}$  are something like infinitely small spherical sectors (with infinitely small aperture and infinitely small radius).

**FixMe:** Describe the co-complete funcoids reverse to these complete funcoids.

THEOREM 2153. A  $D^1$  curve  $f$  is an reparametrized integral curve for a direction field  $d$  iff  $f \in C(\iota_D | \mathbb{R}|_>, Q_+) \cap C(\iota_D | \mathbb{R}|_<, Q_-)$ .

PROOF. Equivalently transform  $f \in C(\iota_D | \mathbb{R}|, Q_+)$ ;  $f \circ \iota_D | \mathbb{R}| \sqsubseteq Q_+ \circ f$ ;  $\langle f \circ \iota_D | \mathbb{R}| \rangle^* @ \{t\} \sqsubseteq \langle Q_+ \circ f \rangle^* @ \{t\}$ ;  $\langle f \rangle^* \Delta_>(t) \sqcap D \sqsubseteq \langle Q_+ \rangle^* f(t)$ ;  $\langle f \rangle^* \Delta_>(t) \sqsubseteq \langle Q_+ \rangle^* f(t)$ ;  $\langle f \rangle^* \Delta_>(t) \sqsubseteq f(t) + W(D(f(t)))$ ;  $\langle f \rangle^* \Delta_>(t) - f(t) \sqsubseteq W(D(f(t)))$ ;

$$\forall r > 0, \phi > 0 \exists \delta > 0 : \langle f \rangle^* (]t; t + \delta[) - f(t) \subseteq R(d(f(t)), \phi) \cap B_r(f(t));$$

$$\forall r > 0, \phi > 0 \exists \delta > 0 \forall 0 < \gamma < \delta : \langle f \rangle^* (]t; t + \gamma[) - f(t) \subseteq R(d(f(t)), \phi) \cap B_r(f(t));$$

$$\forall r > 0, \phi > 0 \exists \delta > 0 \forall 0 < \gamma < \delta : \frac{\langle f \rangle^* (]t; t + \gamma[) - f(t)}{\gamma} \subseteq R(d(f(t)), \phi) \cap B_{r/\delta}(f(t));$$

$$\forall r > 0, \phi > 0 \exists \delta > 0 : \partial_+ f(t) \subseteq R(d(f(t)), \phi) \cap B_{r/\delta}(f(t));$$

$$\forall r > 0, \phi > 0 : \partial_+ f(t) \subseteq R(d(f(t)), \phi);$$

$$\partial_+ f(t) \uparrow\uparrow d(f(t))$$

where  $\partial_+$  is the right derivative.

In the same way we derive that  $C(| \mathbb{R}|_<, Q_-) \Leftrightarrow \partial_- f(t) \uparrow\uparrow d(f(t))$ .

Thus  $f'(t) \uparrow\uparrow d(f(t))$  iff  $f \in C(| \mathbb{R}|_>, Q_+) \cap C(| \mathbb{R}|_<, Q_-)$ .  $\square$

**6.4.  $C^n$  curves.** We consider the differential equation  $f'(t) = d(f(t))$ .

We can consider this equation in any topological vector space  $V$  ([https://en.wikipedia.org/wiki/Frechet\\_derivative](https://en.wikipedia.org/wiki/Frechet_derivative)), see also <https://math.stackexchange.com/q/2977274/4876>. Note that I am not an expert in topological vector spaces and thus my naive generalizations may be wrong in details.

$n$ -th derivative  $f^{(n)}(t) = d_n(f(t))$ ;  $f^{(n+1)}(t) = d'_n(f(t)) \circ f'(t) = d'_n(f(t)) \circ d(f(t))$ . So  $d_{n+1}(y) = d'_n(y) \circ d(y)$ .

Given a point  $y \in V$  define

$$R^n(y) = \left\{ \frac{v \in V}{\overline{vd_0(y) < \frac{d_1(y)}{1!}|v| + \frac{d_2(y)}{2!}|v|^2 + \dots + \frac{d_{n-1}(y)}{(n-1)!}|v|^{n-1} + O(|v|^n), v \neq 0}} \right\}$$

for  $d_0(y) \neq 0$  and  $R^n = \{0\}$  if  $d_0(y) = 0$ .

DEFINITION 2154.  $R^\infty(y) = R^0(y) \sqcap R^1(y) \sqcap R^2(y) \sqcap \dots$

FiXme: It does not work: <https://math.stackexchange.com/a/2978532/4876>.

DEFINITION 2155.  $W^n(y) = R^n(y) \sqcap \prod_{r>0}^{\text{RLD}} B_r(0)$ ;  $W^\infty(y) = R^\infty(y) \sqcap \prod_{r>0}^{\text{RLD}} B_r(0)$ .

Finally our funcoinds are the complete funcoinds  $Q_+^n$  and  $Q_-^n$  described by the formulas

$$\langle Q_+^n \rangle^* @ \{p\} = \langle p+ \rangle W^n(p) \quad \text{and} \quad \langle Q_-^n \rangle^* @ \{p\} = \langle p+ \rangle W^{-n}(p)$$

where  $W^-$  is  $W$  for the reverse vector field  $-d(y)$ .

FiXme: Related questions: <http://math.stackexchange.com/q/1884856/4876> <http://math.stackexchange.com/q/107460/4876> <http://mathoverflow.net/q/88501>

LEMMA 2156. Let for every  $x$  in the domain of the path for small  $t > 0$  we have  $f(x+t) \in R^n(F(f(x)))$  and  $f(x-t) \in R^n(-F(f(x)))$ . Then  $f$  is  $C^n$  smooth.

PROOF. FiXme: Not yet proved!

See also <http://math.stackexchange.com/q/1884930/4876>. □

CONJECTURE 2157. A path  $f$  is conforming to the above differentiable equation and  $C^n$  (where  $n$  is natural or infinite) smooth iff  $f \in C(\iota_D | \mathbb{R}|_{>}, Q_+^n) \cap C(\iota_D | \mathbb{R}|_{<}, Q_-^n)$ .

PROOF. Reverse implication follows from the lemma.

Let now a path  $f$  is  $C^n$ . Then

$$f(x+t) = \sum_{i=0}^n f^{(i)}(x) \frac{t^i}{i!} + o(t^i) = f(x) + f'(x)t + \sum_{i=2}^n f^{(i)}(x) \frac{t^i}{i!} + o(t^i)$$

□

**6.5. Plural funcoinds.** Take  $I_+$  and  $Q_+$  as described above in forward direction and  $I_-$  and  $Q_-$  in backward direction. Then

$$f \in C(I_+, Q_+) \wedge f \in C(I_-, Q_-) \Leftrightarrow f \times f \in C(I_+ \times^{(A)} I_-, Q_+ \times^{(A)} Q_-)?$$

To describe the above we can introduce new term *plural funcoinds*. This is simply a map from an index set to funcoinds. Composition is defined component-wise. Order is defined as product order. Well, do we need this? Isn't it the same as infimum product of funcoinds?



**6.6. Multiple allowed directions per point.**

$$\langle Q \rangle^* @ \{p\} = \bigsqcup_{d \in d(p)} \langle p+ \rangle W(d).$$

It seems (check!) that solutions not only of differential equations but also of difference equations can be expressed as paths in functors.

## CHAPTER 3

### Covers

Let  $S$  be a set of filters.

The corresponding funcoïd (check that it is funcoïd??) is defined as  $\langle f \rangle a = \prod_{\mathcal{X} \in S, a \neq \mathcal{X}} \mathcal{X}$  for atomic filters  $a$ .

A whole defined way to transform a cover into a a funcoïd:

$$f = \bigsqcup_{\mathcal{X} \in S, a \in \text{atoms } \mathcal{X}} (a \times^{(\text{FCD})} a) = \bigsqcup_{\mathcal{X} \in S, \mathcal{C} \sqsubseteq \mathcal{X}} (\mathcal{C} \times^{(\text{FCD})} \mathcal{C}).$$

Is it possible  $\langle f \rangle a \neq \prod_{\mathcal{X} \in S, a \neq \mathcal{X}} \mathcal{X}$ ?

## More on generalized limit

DEFINITION 2158. I will call a permutation group *fixed point free* when every element of it except of identity has no fixed points.

DEFINITION 2159. A funcoid  $f$  is *Kolmogorov* when  $\langle f \rangle^* \{x\} \neq \langle f \rangle^* \{y\}$  for every distinct points  $x, y \in \text{dom } f$ .

### 1. Hausdorff funcoids

DEFINITION 2160. *Limit*  $\lim \mathcal{F} = x$  of a filter  $\mathcal{F}$  regarding funcoid  $f$  is such a point that  $\langle f \rangle^* \{x\} \supseteq \mathcal{F}$ .

DEFINITION 2161. *Hausdorff* funcoid is such a funcoid that every proper filter on its image has at most one limit.

PROPOSITION 2162. The following are pairwise equivalent for every funcoid  $f$ :

- 1°.  $f$  is Hausdorff.
- 2°.  $x \neq y \Rightarrow \langle f \rangle^* \{x\} \asymp \langle f \rangle^* \{y\}$ .

PROOF.

1°  $\Rightarrow$  2°. If 2° does not hold, then there exist distinct points  $x$  and  $y$  such that  $\langle f \rangle^* \{x\} \not\asymp \langle f \rangle^* \{y\}$ . So  $x$  and  $y$  are both limit points of  $\langle f \rangle^* \{x\} \cap \langle f \rangle^* \{y\}$ , and thus  $f$  is not Hausdorff.

2°  $\Rightarrow$  1°. Suppose  $\mathcal{F}$  is proper.

$$\langle f \rangle^* \{x\} \supseteq \mathcal{F} \wedge \langle f \rangle^* \{y\} \supseteq \mathcal{F} \Rightarrow \langle f \rangle^* \{x\} \not\asymp \langle f \rangle^* \{y\} \Rightarrow x = y.$$

□

COROLLARY 2163. Every entirely defined Hausdorff funcoid is Kolmogorov.

REMARK 2164. It is enough to be “almost entirely defined” (having nonempty value everywhere except of one point).

OBVIOUS 2165. For a complete funcoid induced by a topological space this coincides with the traditional definition of a Hausdorff topological space.

### 2. Restoring functions from limit

Consider alternative definition of generalized limit:

$$\text{xlim } f = \lambda r \in G : \nu \circ f \circ \uparrow r.$$

Or:

$$\text{xlim}_a f = \left\{ \frac{(\langle r^{-1} \rangle^* a, \nu \circ f \circ \uparrow r)}{r \in G} \right\}$$

(note this requires explicit filter in the definition of generalized limit).

Operations on the set of generalized limits can be defined (twice) pointwise.

**FiXme:** First define operations on funcoids.

PROPOSITION 2166. The above defined  $\text{xlim}_{\langle \mu \rangle^* \{x\}} f$  is a monovalued function if  $\mu$  is Kolmogorov and  $G$  is fixed point free.

PROOF. We need to prove  $\langle r^{-1} \rangle \langle \mu \rangle^* \{x\} \neq \langle s^{-1} \rangle \langle \mu \rangle^* \{x\}$  for  $r, s \in G$ ,  $r \neq s$ . Really, by definition of generalized limit, they commute, so our formula is equivalent to  $\langle \mu \rangle^* \langle r^{-1} \rangle^* \{x\} \neq \langle \mu \rangle^* \langle s^{-1} \rangle^* \{x\}$ ;  $\langle \mu \rangle^* \langle r^{-1} \circ s \rangle^* \langle s^{-1} \rangle^* \{x\} \neq \langle \mu \rangle^* \langle s^{-1} \rangle^* \{x\}$ . But  $r^{-1} \circ s \neq e$ , so because it is fixed point free,  $\langle r^{-1} \circ s \rangle^* \langle s^{-1} \rangle^* \{x\} \neq \langle s^{-1} \rangle^* \{x\}$  and thus by kolmogorovness, we have the thesis.  $\square$

LEMMA 2167. Let  $\mu$  and  $\nu$  be Hausdorff funcoids. If function  $f$  is defined at point  $x$ , then

$$fx = \lim \langle (\text{xlim}_{\langle \mu \rangle^* \{x\}} f) \langle \mu \rangle^* \{x\} \rangle^* \{x\}$$

REMARK 2168. The right part is correctly defined because  $\text{xlim}_a f$  is monoalued.

PROOF.  $\lim \langle (\text{xlim}_{\langle \mu \rangle^* \{x\}} f) \langle \mu \rangle^* \{x\} \rangle^* \{x\} = \lim \langle \nu \circ f \rangle^* \{x\} = \lim \langle \nu \rangle^* fx = fx$ .  $\square$

COROLLARY 2169. Let  $\mu$  and  $\nu$  be Hausdorff funcoids. Then function  $f$  can be restored from values of  $\text{xlim}_{\langle \mu \rangle^* \{x\}} f$ .

## Extending Galois connections between functors and relocks

DEFINITION 2170.

- 1°.  $\Phi_* f = \lambda b \in \mathfrak{B} : \sqcup \left\{ \frac{x \in \mathfrak{A}}{fx \sqsubseteq b} \right\};$   
 2°.  $\Phi^* f = \lambda b \in \mathfrak{A} : \prod \left\{ \frac{x \in \mathfrak{B}}{fx \sqsupseteq b} \right\}.$

PROPOSITION 2171.

- 1°. If  $f$  has upper adjoint then  $\Phi_* f$  is the upper adjoint of  $f$ .  
 2°. If  $f$  has lower adjoint then  $\Phi^* f$  is the lower adjoint of  $f$ .

PROOF. By theorem 131. □

LEMMA 2172.  $\Phi^*(\text{RLD})_{\text{out}} = (\text{FCD}).$

PROOF.  $(\Phi^*(\text{RLD})_{\text{out}})f = \prod \left\{ \frac{g \in \text{FCD}}{(\text{RLD})_{\text{out}} g \sqsupseteq f} \right\} = \prod^{\text{FCD}} \left\{ \frac{g \in \mathbf{Rel}}{(\text{RLD})_{\text{out}} g \sqsupseteq f} \right\} =$   
 $\prod^{\text{FCD}} \left\{ \frac{g \in \mathbf{Rel}}{g \sqsupseteq f} \right\} = (\text{FCD})f.$  □

LEMMA 2173.  $\Phi_*(\text{RLD})_{\text{out}} \neq (\text{FCD}).$

PROOF.  $(\Phi_*(\text{RLD})_{\text{out}})f = \sqcup \left\{ \frac{g \in \text{FCD}}{(\text{RLD})_{\text{out}} g \sqsubseteq f} \right\}$   
 $(\Phi_*(\text{RLD})_{\text{out}}) \perp \neq \perp.$  □

LEMMA 2174.  $\Phi^*(\text{FCD}) = (\text{RLD})_{\text{out}}.$

PROOF.  $(\Phi^*(\text{FCD}))f = \prod \left\{ \frac{g \in \text{RLD}}{(\text{FCD})g \sqsupseteq f} \right\} = \prod^{\text{RLD}} \left\{ \frac{g \in \mathbf{Rel}}{(\text{FCD})g \sqsupseteq f} \right\} = \prod^{\text{RLD}} \left\{ \frac{g \in \mathbf{Rel}}{g \sqsupseteq f} \right\} =$   
 $(\text{RLD})_{\text{out}}f.$  □

LEMMA 2175.  $\Phi_*(\text{RLD})_{\text{in}} = (\text{FCD}).$

PROOF.  $(\Phi_*(\text{RLD})_{\text{in}})f = \sqcup \left\{ \frac{g \in \text{FCD}}{(\text{RLD})_{\text{in}} g \sqsubseteq f} \right\} = \sqcup \left\{ \frac{g \in \text{FCD}}{g \sqsubseteq (\text{FCD})f} \right\} = (\text{FCD})f.$  □

THEOREM 2176. The picture at figure 14 describes values of functions  $\Phi_*$  and  $\Phi^*$ . All nodes of this diagram are distinct.

PROOF. Follows from the above lemmas. □




FIGURE 14

QUESTION 2177. What is at the node “other”?

Trying to answer this question:

LEMMA 2178.  $(\Phi_*(\text{RLD})_{\text{out}})\perp = \Omega^{\text{FCD}}$ .

PROOF. We have  $(\text{RLD})_{\text{out}}\Omega^{\text{FCD}} = \perp$ .  $x \not\sqsubseteq \Omega^{\text{FCD}} \Rightarrow (\text{RLD})_{\text{out}}x \sqsupseteq \text{Cor } x \sqsupset \perp$ .

Thus  $\max\left\{\frac{x \in \text{FCD}}{(\text{RLD})_{\text{out}}x = \perp}\right\} = \Omega^{\text{FCD}}$ .

So  $(\Phi_*(\text{RLD})_{\text{out}})\perp = \Omega^{\text{FCD}}$ . □

CONJECTURE 2179.  $(\Phi_*(\text{RLD})_{\text{out}})f = \Omega^{\text{FCD}} \sqcup (\text{FCD})f$ .

The above conjecture looks not natural, but I do not see a better alternative formula.

QUESTION 2180. What happens if we keep applying  $\Phi^*$  and  $\Phi_*$  to the node “other”? Will we this way get a finite or infinite set?

## Boolean funcoids

### 1. One-element boolean lattice

Let  $\mathfrak{A}$  be a boolean lattice and  $\mathfrak{B} = \mathcal{P}0$ . It's sole element is  $\perp$ .

$$f \in \text{pFCD}(\mathfrak{A}; \mathfrak{B}) \Leftrightarrow \forall X \in \mathfrak{A} : (\langle f \rangle X \neq \perp \Leftrightarrow \langle f^{-1} \rangle \perp \neq X) \Leftrightarrow \forall X \in \mathfrak{A} : (0 \Leftrightarrow \langle f^{-1} \rangle \perp \neq X) \Leftrightarrow \forall X \in \mathfrak{A} : \langle f^{-1} \rangle \perp \simeq X \Leftrightarrow \forall X \in \mathfrak{A} : \langle f^{-1} \rangle \perp = \perp^{\mathfrak{A}} \Leftrightarrow \langle f^{-1} \rangle \perp = \perp^{\mathfrak{A}} \Leftrightarrow \langle f^{-1} \rangle = \{(\perp; \perp^{\mathfrak{A}})\}.$$

Thus  $\text{card pFCD}(\mathfrak{A}; \mathcal{P}0) = 1$ .

### 2. Two-element boolean lattice

Consider the two-element boolean lattice  $\mathfrak{B} = \mathcal{P}1$ .

Let  $f$  be a pointfree protofuncoid from  $\mathfrak{A}$  to  $\mathfrak{B}$  (that is  $(\mathfrak{A}; \mathfrak{B}; \alpha; \beta)$  where  $\alpha \in \mathfrak{B}^{\mathfrak{A}}, \beta \in \mathfrak{A}^{\mathfrak{B}}$ ).

$$f \in \text{pFCD}(\mathfrak{A}; \mathfrak{B}) \Leftrightarrow \forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\langle f \rangle X \neq Y \Leftrightarrow \langle f^{-1} \rangle Y \neq X) \Leftrightarrow \forall X \in \mathfrak{A}, Y \in \mathfrak{B} : ((0 \in \langle f \rangle X \wedge 0 \in Y) \vee (1 \in \langle f \rangle X \wedge 1 \in Y) \Leftrightarrow \langle f^{-1} \rangle Y \neq X).$$

$T = \left\{ \frac{X \in \mathfrak{A}}{0 \in \langle f \rangle X} \right\}$  is an ideal. Really: That it's an upper set is obvious. Let  $P \cup Q \in \left\{ \frac{X \in \mathfrak{A}}{0 \in \langle f \rangle X} \right\}$ . Then  $0 \in \langle f \rangle (P \cup Q) = \langle f \rangle P \cup \langle f \rangle Q$ ;  $0 \in \langle f \rangle P \vee 0 \in \langle f \rangle Q$ .

Similarly  $S = \left\{ \frac{X \in \mathfrak{A}}{1 \in \langle f \rangle X} \right\}$  is an ideal.

Let now  $T, S \in \mathcal{P}\mathfrak{A}$  be ideals. Can we restore  $\langle f \rangle$ ? Yes, because we know  $0 \in \langle f \rangle X$  and  $1 \in \langle f \rangle X$  for every  $X \in \mathfrak{A}$ .

So it is equivalent to  $\forall X \in \mathfrak{A}, Y \in \mathfrak{B} : ((X \in T \wedge 0 \in Y) \vee (X \in S \wedge 1 \in Y) \Leftrightarrow \langle f^{-1} \rangle Y \neq X)$ .

$f \in \text{pFCD}(\mathfrak{A}; \mathfrak{B})$  is equivalent to conjunction of all rows of this table:

$Y$	equality
$\emptyset$	$\langle f^{-1} \rangle \emptyset = \emptyset$
$\{0\}$	$X \in T \Leftrightarrow \langle f^{-1} \rangle \{0\} \neq X$
$\{1\}$	$X \in S \Leftrightarrow \langle f^{-1} \rangle \{1\} \neq X$
$\{0,1\}$	$X \in T \vee X \in S \Leftrightarrow \langle f^{-1} \rangle \{0,1\} \neq X$

Simplified:

$Y$	equality
$\emptyset$	$\langle f^{-1} \rangle \emptyset = \emptyset$
$\{0\}$	$T = \partial \langle f^{-1} \rangle \{0\}$
$\{1\}$	$S = \partial \langle f^{-1} \rangle \{1\}$
$\{0,1\}$	$T \cup S = \partial \langle f^{-1} \rangle \{0,1\}$

From the last table it follows that  $T$  and  $S$  are principal ideals.

So we can take arbitrary either  $\langle f^{-1} \rangle \{0\}$ ,  $\langle f^{-1} \rangle \{1\}$  or principal ideals  $T$  and  $S$ .

In other words, we take  $\langle f^{-1} \rangle \{0\}$ ,  $\langle f^{-1} \rangle \{1\}$  arbitrary and independently. So we have  $\text{pFCD}(\mathfrak{A}; \mathfrak{B})$  equivalent to product of two instances of  $\mathfrak{A}$ . So it a boolean lattice. **FiXme: I messed product with disjoint union below.)**

### 3. Finite boolean lattices

We can assume  $\mathfrak{B} = \mathcal{P}B$  for a set  $B$ ,  $\text{card } B = n$ . Then

$$f \in \text{pFCD}(\mathfrak{A}; \mathfrak{B}) \Leftrightarrow \forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\langle f \rangle X \neq Y \Leftrightarrow \langle f^{-1} \rangle Y \neq X) \Leftrightarrow \forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\exists i \in Y : i \in \langle f \rangle X \Leftrightarrow \langle f^{-1} \rangle Y \neq X).$$

Having values of  $\langle f^{-1} \rangle \{i\}$  we can restore all  $\langle f^{-1} \rangle Y$ . [need this paragraph?]

$$\text{Let } T_i = \left\{ \frac{X \in \mathfrak{A}}{i \in \langle f \rangle X} \right\}.$$

Let now  $T_i \in \mathcal{P}\mathfrak{A}$  be ideals. Can we restore  $\langle f \rangle$ ? Yes, because we know  $i \in \langle f \rangle X$  for every  $X \in \mathfrak{A}$ .

So, it is equivalent to:

$$\forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\exists i \in Y : X \in T_i \Leftrightarrow \langle f^{-1} \rangle Y \neq X). \quad (1)$$

LEMMA 2181. The formula (1) is equivalent to:

$$\forall X \in \mathfrak{A}, i \in B : (X \in T_i \Leftrightarrow \langle f^{-1} \rangle \{i\} \neq X). \quad (2)$$

PROOF. (1) $\Rightarrow$ (2). Just take  $Y = \{i\}$ .

(2) $\Rightarrow$ (1). Let (2) holds. Let also  $X \in \mathfrak{A}, Y \in \mathfrak{B}$ . Then  $\langle f^{-1} \rangle Y \neq X \Leftrightarrow \bigcup_{i \in Y} \langle f^{-1} \rangle \{i\} \neq X \Leftrightarrow \exists i \in Y : \langle f^{-1} \rangle \{i\} \neq X \Leftrightarrow \exists i \in Y : X \in T_i$ .  $\square$

Further transforming:  $\forall i \in B : T_i = \partial \langle f^{-1} \rangle \{i\}$ .

So  $\langle f^{-1} \rangle \{i\}$  are arbitrary elements of  $\mathfrak{B}$  and  $T_i$  are corresponding arbitrary principal ideals.

In other words,  $\text{pFCD}(\mathfrak{A}; \mathfrak{B}) \cong \mathfrak{A} \Pi \dots \Pi \mathfrak{A}$  ( $\text{card } B$  times). Thus  $\text{pFCD}(\mathfrak{A}; \mathfrak{B})$  is a boolean lattice.

### 4. About infinite case

Let  $\mathfrak{A}$  be a complete boolean lattice,  $\mathfrak{B}$  be an atomistic boolean lattice.

$$f \in \text{pFCD}(\mathfrak{A}; \mathfrak{B}) \Leftrightarrow \forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\langle f \rangle X \neq Y \Leftrightarrow \langle f^{-1} \rangle Y \neq X) \Leftrightarrow \forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\exists i \in \text{atoms } Y : i \in \text{atoms } \langle f \rangle X \Leftrightarrow \langle f^{-1} \rangle Y \neq X).$$

$$\text{Let } T_i = \left\{ \frac{X \in \mathfrak{A}}{i \in \text{atoms } \langle f \rangle X} \right\}.$$

$T_i$  is an ideal: Really: That it's an upper set is obvious. Let  $P \cup Q \in \left\{ \frac{X \in \mathfrak{A}}{i \in \text{atoms } \langle f \rangle X} \right\}$ . Then  $i \in \text{atoms } \langle f \rangle (P \cup Q) = \text{atoms } \langle f \rangle P \cup \text{atoms } \langle f \rangle Q; i \in \langle f \rangle P \vee i \in \langle f \rangle Q$ .

Let now  $T_i \in \mathcal{P}\mathfrak{A}$  be ideals. Can we restore  $\langle f \rangle$ ? Yes, because we know  $i \in \text{atoms } \langle f \rangle X$  for every  $X \in \mathfrak{A}$  and  $\mathfrak{B}$  is atomistic.

So, it is equivalent to:

$$\forall X \in \mathfrak{A}, Y \in \mathfrak{B} : (\exists i \in \text{atoms } Y : X \in T_i \Leftrightarrow \langle f^{-1} \rangle Y \neq X). \quad (3)$$

LEMMA 2182. The formula (3) is equivalent to:

$$\forall X \in \mathfrak{A}, i \in \text{atoms } \mathfrak{B} : (X \in T_i \Leftrightarrow \langle f^{-1} \rangle i \neq X). \quad (4)$$

PROOF. (3) $\Rightarrow$ (4). Let (3) holds. Take  $Y = i$ . Then  $\text{atoms } Y = \{i\}$  and thus  $X \in T_i \Leftrightarrow \exists i \in \text{atoms } Y : X \in T_i \Leftrightarrow \langle f^{-1} \rangle Y \neq X \Leftrightarrow \langle f^{-1} \rangle i \neq X$ .

(4) $\Rightarrow$ (3). Let (4) holds. Let also  $X \in \mathfrak{A}, Y \in \mathfrak{B}$ . Then  $\langle f^{-1} \rangle Y \neq X \Leftrightarrow \langle f^{-1} \rangle \bigsqcup \text{atoms } Y \neq X \Leftrightarrow \bigsqcup_{i \in \text{atoms } Y} \langle f^{-1} \rangle i \neq X \Leftrightarrow \exists i \in \text{atoms } Y : \langle f^{-1} \rangle i \neq X \Leftrightarrow \exists i \in \text{atoms } Y : X \in T_i$ .  $\square$

Further equivalently transforming:  $\forall i \in \text{atoms } \mathfrak{B} : T_i = \partial \langle f^{-1} \rangle i$ .

So  $\langle f^{-1} \rangle i$  are arbitrary elements of  $\mathfrak{B}$  and  $T_i$  are corresponding arbitrary principal ideals.



In other words,  $\text{pFCD}(\mathfrak{A}; \mathfrak{B}) \cong \prod_{i \in \text{card atoms } \mathfrak{B}} \mathfrak{A}$ . Thus  $\text{pFCD}(\mathfrak{A}; \mathfrak{B})$  is a boolean lattice.

So finally we have a very weird theorem, which is a partial solution for the above open problem (The weirdness is in its partiality and asymmetry):

**THEOREM 2183.** If  $\mathfrak{A}$  is a complete boolean lattice and  $\mathfrak{B}$  is an atomistic boolean lattice (or vice versa), then  $\text{pFCD}(\mathfrak{A}; \mathfrak{B})$  is a boolean lattice.

[4] proves “THEOREM 4.6. Let  $A, B$  be bounded posets.  $A \otimes B$  is a completely distributive complete Boolean lattice iff  $A$  and  $B$  are completely distributive Boolean lattices.” (where  $A \otimes B$  is equivalent to the set of Galois connections between  $A$  and  $B$ ) and other interesting results.

## Interior funcoids

Having a funcoid  $f$  let define *interior funcoid*  $f^\circ$ .

DEFINITION 2184. Let  $f \in \text{FCD}(A, B) = \text{pFCD}(\mathcal{T}A, \mathcal{T}B)$  be a co-complete funcoid. Then  $f^\circ \in \text{pFCD}(\text{dual } \mathcal{T}A, \text{dual } \mathcal{T}B)$  is defined by the formula  $\langle f^\circ \rangle^* X = \overline{\langle f \rangle X}$ .

PROPOSITION 2185. Pointfree funcoid  $f^\circ$  exists and is unique.

PROOF.  $X \mapsto \overline{\langle f \rangle X}$  is a component of pointfree funcoid  $\text{dual } \mathcal{T}A \rightarrow \text{dual } \mathcal{T}B$  iff  $\langle f \rangle$  is a component of the corresponding pointfree funcoid  $\mathcal{T}A \rightarrow \mathcal{T}B$  that is essentially component of the corresponding funcoid  $\text{FCD}(A, B)$  what holds for a unique funcoid.  $\square$

It can be also defined for arbitrary funcoids by the formula  $f^\circ = (\text{CoCompl } f)^\circ$ .

OBVIOUS 2186.  $f^\circ$  is co-complete.

THEOREM 2187. The following values are pairwise equal for a co-complete funcoid  $f$  and  $X \in \mathcal{T} \text{Src } f$ :

- 1 $^\circ$ .  $\langle f^\circ \rangle^* X$ ;
- 2 $^\circ$ .  $\left\{ \frac{y \in \text{Dst } f}{\langle f^{-1} \rangle^* \{y\} \sqsubseteq X} \right\}$
- 3 $^\circ$ .  $\bigsqcup \left\{ \frac{Y \in \mathcal{T} \text{Dst } f}{\langle f^{-1} \rangle^* Y \sqsubseteq X} \right\}$
- 4 $^\circ$ .  $\bigsqcup \left\{ \frac{\mathcal{Y} \in \mathcal{F} \text{Dst } f}{\langle f^{-1} \rangle \mathcal{Y} \sqsubseteq X} \right\}$

PROOF.

$$1^\circ = 2^\circ. \left\{ \frac{y \in \text{Dst } f}{\langle f^{-1} \rangle^* \{y\} \sqsubseteq X} \right\} = \left\{ \frac{x \in \text{Dst } f}{\langle f^{-1} \rangle^* \{x\} \succ \overline{X}} \right\} = \left\{ \frac{x \in \text{Dst } f}{\{x\} \succ \langle f \rangle \overline{X}} \right\} = \overline{\langle f \rangle \overline{X}} = \langle f^\circ \rangle^* X.$$

2 $^\circ$  = 3 $^\circ$ . If  $\langle f^{-1} \rangle^* Y \sqsubseteq X$  then (by completeness of  $f^{-1}$ )  $Y = \left\{ \frac{y \in Y}{\langle f^{-1} \rangle^* \{y\} \sqsubseteq X} \right\}$  and thus

$$\bigsqcup \left\{ \frac{Y \in \mathcal{T} \text{Dst } f}{\langle f^{-1} \rangle^* Y \sqsubseteq X} \right\} \sqsubseteq \left\{ \frac{y \in \text{Dst } f}{\langle f^{-1} \rangle^* \{y\} \sqsubseteq X} \right\}.$$

The reverse inequality is obvious.

3 $^\circ$  = 4 $^\circ$ . It's enough to prove that if  $\langle f^{-1} \rangle \mathcal{Y} \sqsubseteq X$  for  $\mathcal{Y} \in \mathcal{F} \text{Dst } f$  then exists  $Y \in \text{up } \mathcal{Y}$  such that  $\langle f^{-1} \rangle^* Y \sqsubseteq X$ . Really let  $\langle f^{-1} \rangle \mathcal{Y} \sqsubseteq X$ . Then  $\bigsqcap \langle \langle f^{-1} \rangle^* \rangle \text{up } \mathcal{Y} \sqsubseteq X$  and thus exists  $Y \in \text{up } \mathcal{Y}$  such that  $\langle f^{-1} \rangle^* Y \sqsubseteq X$  by properties of generalized filter bases.  $\square$

This coincides with the customary definition of interior in topological spaces.

PROPOSITION 2188.  $f^{\circ\circ} = f$  for every funcoid  $f$ .

PROOF.  $\langle f^{\circ\circ} \rangle^* X = \neg \neg \langle f \rangle \neg \neg X = \langle f \rangle X$ .  $\square$

PROPOSITION 2189. Let  $g \in \text{FCD}(A, B)$ ,  $f \in \text{FCD}(B, C)$ ,  $h \in \text{FCD}(A, C)$  for some sets  $A, B, C$ .

$g \sqsubseteq f^\circ \circ h \Leftrightarrow f^{-1} \circ g \sqsubseteq h$ , provided  $f$  and  $h$  are co-complete.

PROOF.  $g \sqsubseteq f^\circ \circ h \Leftrightarrow \forall X \in A : \langle g \rangle^* X \sqsubseteq \langle f^\circ \circ h \rangle^* X \Leftrightarrow \forall X \in A : \langle g \rangle^* X \sqsubseteq \langle f^\circ \rangle^* \langle h \rangle^* X \Leftrightarrow \forall X \in A : \langle g \rangle^* X \sqsubseteq \neg \langle f \rangle^* \neg \langle h \rangle^* X \Leftrightarrow \forall X \in A : \langle g \rangle^* X \simeq \langle f \rangle^* \neg \langle h \rangle^* X \Leftrightarrow \forall X \in A : \langle f^{-1} \rangle^* \langle g \rangle^* X \simeq \neg \langle h \rangle^* X \Leftrightarrow \forall X \in A : \langle f^{-1} \rangle^* \langle g \rangle^* X \sqsubseteq \langle h \rangle^* X \Leftrightarrow \forall X \in A : \langle f^{-1} \circ g \rangle^* X \sqsubseteq \langle h \rangle^* X \Leftrightarrow f^{-1} \circ g \sqsubseteq h. \quad \square$

REMARK 2190. The above theorem allows to get rid of interior functors (and use only “regular” functors) in some formulas.

## Filterization of pointfree funcoids

Let  $(\mathfrak{A}, \mathfrak{Z}_0)$  and  $(\mathfrak{B}, \mathfrak{Z}_1)$  be primary filtrators over boolean lattices. By corollary 518 we have that  $\mathfrak{A}$  and  $\mathfrak{B}$  are complete lattices.

Let  $f$  be a pointfree funcoid  $\mathfrak{Z}_0 \rightarrow \mathfrak{Z}_1$ . Define pointfree funcoid  $\uparrow f$  (*filterization* of  $f$ ) by the formulas

$$\langle \uparrow f \rangle \mathcal{X} = \prod_{X \in \text{up } \mathcal{X}}^{\mathfrak{B}} \langle f \rangle X \quad \text{and} \quad \langle \uparrow f^{-1} \rangle \mathcal{Y} = \prod_{Y \in \text{up } \mathcal{Y}}^{\mathfrak{A}} \langle f^{-1} \rangle Y.$$

PROPOSITION 2191.  $\uparrow f$  is a pointfree funcoid.

PROOF.

$$\begin{aligned} \mathcal{Y} \neq \langle \uparrow f \rangle \mathcal{X} &\Leftrightarrow \mathcal{Y} \neq \prod_{X \in \text{up } \mathcal{X}}^{\mathfrak{B}} \langle f \rangle X \Leftrightarrow \\ &\prod_{X \in \text{up } \mathcal{X}}^{\mathfrak{B}} (\mathcal{Y} \cap^{\mathfrak{B}} \langle f \rangle X) \neq \perp \Leftrightarrow \text{(corollary 573*)} \\ &\forall X \in \text{up } \mathcal{X} : \mathcal{Y} \cap^{\mathfrak{B}} \langle f \rangle X \neq \perp \Leftrightarrow \text{(theorem 537)} \\ &\forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y} : Y \cap^{\mathfrak{B}} \langle f \rangle X \neq \perp \Leftrightarrow \text{(corollary 536)} \\ &\forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y} : Y \cap^{\mathfrak{Z}_1} \langle f \rangle X \neq \perp \Leftrightarrow \\ &\forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y} : X [f] Y. \end{aligned}$$

\* To apply corollary 573 we need to show that  $\left\{ \frac{\mathcal{Y} \cap^{\mathfrak{B}} \langle f \rangle X}{X \in \text{up } \mathcal{X}} \right\}$  is a generalized filter base. To show it is enough to show that  $\left\{ \frac{\langle f \rangle X}{X \in \text{up } \mathcal{X}} \right\}$  is a generalized filter base. But this easily follows from proposition 1603 and 579.

Similarly  $\mathcal{X} \neq \langle \uparrow f^{-1} \rangle \mathcal{Y} \Leftrightarrow \forall X \in \text{up } \mathcal{X}, Y \in \text{up } \mathcal{Y} : X [f] Y$ . Thus  $\mathcal{Y} \neq \langle \uparrow f \rangle \mathcal{X} \Leftrightarrow \mathcal{X} \neq \langle \uparrow f^{-1} \rangle \mathcal{Y}$ .  $\square$

PROPOSITION 2192. The above defined  $\uparrow$  is an injection.

PROOF.  $\langle \uparrow f \rangle X = \prod_{X' \in \text{up } X}^{\mathfrak{B}} \langle f \rangle X' = \min_{X' \in \text{up } X} \langle f \rangle X' = \langle f \rangle X$ . So  $\langle f \rangle$  is determined by  $\langle \uparrow f \rangle$ . Likewise  $\langle f^{-1} \rangle$  is determined by  $\langle \uparrow f^{-1} \rangle$ .  $\square$

CONJECTURE 2193. (Non generalizing of theorem 1712) Pointfree funcoids  $f$  between some: a. atomistic but non-complete; b. complete but non-atomistic boolean lattices  $\mathfrak{Z}_0$  and  $\mathfrak{Z}_1$  do not bijectively correspond to morphisms  $p \in \mathbf{Rel}(\text{atoms } \mathfrak{Z}_0, \text{atoms } \mathfrak{Z}_1)$  by the formulas:

$$\begin{aligned} \langle f \rangle X &= \bigsqcup \langle p \rangle^* \text{atoms } X, \quad \langle f^{-1} \rangle Y = \bigsqcup \langle p^{-1} \rangle^* \text{atoms } Y; \\ (x, y) \in \text{GR } p &\Leftrightarrow y \in \text{atoms } \langle f \rangle x \Leftrightarrow x \in \text{atoms } \langle f^{-1} \rangle y. \end{aligned}$$

## Systems of sides

Now we will consider a common generalization of (some of pointfree) functors and (some of) Galois connections. The main purpose of this is general theorem 2241 below.

First consider some properties of Galois connections:

### 1. More on Galois connections

Here I will denote  $\langle f \rangle$  the lower adjoint of a Galois connection  $f$ . **FiXme:** Switch to this notation in the book?

Let  $\mathbf{GAL}$  be the category of Galois connections. **FiXme:** Need to decide whether use  $\mathbf{GAL}(A, B)$  or  $A \otimes B$ .

I will denote  $(f, g)^{-1} = (g, f)$  for a Galois connection  $(f, g)$ .

We will order Galois connections by the formula

$$f \sqsubseteq g \Leftrightarrow \langle f \rangle \sqsubseteq \langle g \rangle \Leftrightarrow \langle f^{-1} \rangle \supseteq \langle g^{-1} \rangle.$$

OBVIOUS 2194. This defines a partial order on the set of Galois connections between any two (fixed) posets.

PROPOSITION 2195. If  $f$  and  $g$  are Galois connections (between a join-semilattice  $\mathfrak{A}$  and a meet-semilattice  $\mathfrak{B}$ ), then there exists a Galois connection  $f \sqcup g$  determined by the formula  $\langle f \sqcup g \rangle x = \langle f \rangle x \sqcup \langle g \rangle x$ .

PROOF. It is enough to prove that

$$(x \mapsto \langle f \rangle x \sqcup \langle g \rangle x, y \mapsto \langle f^{-1} \rangle y \sqcap \langle g^{-1} \rangle y)$$

is a Galois connection that is that

$$\langle f \rangle x \sqcup \langle g \rangle x \sqsubseteq y \Leftrightarrow x \sqsubseteq \langle f^{-1} \rangle y \sqcap \langle g^{-1} \rangle y$$

for all relevant  $x$  and  $y$ .

Really,

$$\begin{aligned} \langle f \rangle x \sqcup \langle g \rangle x \sqsubseteq y &\Leftrightarrow \langle f \rangle x \sqsubseteq y \wedge \langle g \rangle x \sqsubseteq y \Leftrightarrow \\ &x \sqsubseteq \langle f^{-1} \rangle y \wedge x \sqsubseteq \langle g^{-1} \rangle y \Leftrightarrow x \sqsubseteq \langle f^{-1} \rangle y \sqcap \langle g^{-1} \rangle y. \end{aligned}$$

□

**FiXme:** Describe infinite join of Galois connections.

PROPOSITION 2196. If  $\mathfrak{A}$  is a poset with least element, then  $\langle a \rangle \perp = \perp$ .

PROOF.  $\langle a \rangle \perp \sqsubseteq y \Leftrightarrow \perp \sqsubseteq \langle a^{-1} \rangle y \Leftrightarrow 1$ . Thus  $\langle a \rangle \perp$  is the least element. □

PROPOSITION 2197.  $(\mathfrak{A} \times \{\perp^{\mathfrak{B}}\}, \mathfrak{B} \times \{\top^{\mathfrak{A}}\})$  is the least Galois connection from a poset  $\mathfrak{A}$  with greatest element to a poset  $\mathfrak{B}$  with least element.

PROOF. Let's prove that it is a Galois connection. We need to prove

$$(\mathfrak{A} \times \{\perp^{\mathfrak{B}}\})x \sqsubseteq y \Leftrightarrow x \sqsubseteq (\mathfrak{B} \times \{\top^{\mathfrak{A}}\})y.$$

But this is trivially equivalent to  $1 \Leftrightarrow 1$ . Thus it's a Galois connection.

That it the least is obvious. □

COROLLARY 2198.  $\langle \perp \rangle x = \perp$  for Galois connections from a poset  $\mathfrak{A}$  with greatest element to a poset  $\mathfrak{B}$  with least element. **FixMe: Clarify.**

THEOREM 2199. If  $\mathfrak{A}$  and  $\mathfrak{B}$  are bounded posets, then  $\text{GAL}(\mathfrak{A}, \mathfrak{B})$  is bounded.

PROOF. That  $\text{GAL}(\mathfrak{A}, \mathfrak{B})$  has least element was proved above. I will demonstrate that  $(\alpha, \beta)$  is the greatest element of  $\text{pFCD}(\mathfrak{A}, \mathfrak{B})$  for

$$\alpha X = \begin{cases} \perp^{\mathfrak{B}} & \text{if } X = \perp^{\mathfrak{A}} \\ \top^{\mathfrak{B}} & \text{if } X \neq \perp^{\mathfrak{A}} \end{cases}; \quad \beta Y = \begin{cases} \top^{\mathfrak{A}} & \text{if } Y = \top^{\mathfrak{B}} \\ \perp^{\mathfrak{A}} & \text{if } Y \neq \top^{\mathfrak{B}} \end{cases}.$$

First prove  $Y \sqsubseteq \alpha X \Leftrightarrow X \sqsubseteq \beta Y$ .

Really  $\alpha X \sqsubseteq Y \Leftrightarrow X = \perp^{\mathfrak{A}} \vee Y = \top^{\mathfrak{B}} \Leftrightarrow X \sqsubseteq \beta Y$ .

That it is the greatest Galois connection between  $\mathfrak{A}$  and  $\mathfrak{B}$  easily follows from proposition 2196.  $\square$

THEOREM 2200. For every brouwerian lattice  $x \mapsto c \sqcap x$  is a lower adjoint.

PROOF. By dual of theorem 154.  $\square$

EXERCISE 2201. Describe the corresponding upper adjoint, especially for the special case of boolean lattices.

## 2. Definition

DEFINITION 2202. *System of presides* is a functor  $\Upsilon = (f \mapsto \langle f \rangle)$  from an ordered category to the category of functions between (small) bounded lattices, such that (for all relevant variables):

- 1°. Every  $\text{Hom}$ -set of  $\text{Src } \Upsilon$  is a bounded join-semilattice.
- 2°.  $\langle a \rangle \perp = \perp$ .
- 3°.  $\langle a \sqcup b \rangle X = \langle a \rangle X \sqcup \langle b \rangle X$  (equivalent to  $\Upsilon$  to be a join-semilattice homomorphism, if we order functions between small bounded lattices component-wise).

I call morphisms of such categories *sides*.<sup>1</sup>

REMARK 2203. We could generalize to functions between small join-semilattices with least elements instead of bounded lattices only, but this is not really necessary.

DEFINITION 2204. I will call objects of the source category of this functor simply *objects of the presides*.

DEFINITION 2205. *Bounded system of presides* is system of presides from an ordered category with bounded  $\text{Hom}$ -sets such that  $X, Y \in \text{Ob Src } \Upsilon$  the following additional axioms hold for all suitable  $a$ :

- 1°.  $\langle \perp^{\text{Hom}(X, Y)} \rangle a = \perp$ .
- 2°.  $\langle \top^{\text{Hom}(X, Y)} \rangle a = \top$  unless  $a = \perp$

DEFINITION 2206. *System of presides with identities* is a system of presides with a morphism  $\text{id}_a \in \text{Src } \Upsilon$  for every object  $\mathfrak{A}$  of  $\text{Src } \Upsilon$  and  $a \in \mathfrak{A}$  and the following additional axioms:

- 1°.  $\text{id}_c \sqsubseteq 1_{\mathfrak{A}}$  for every  $c \in \mathfrak{A}$  where  $\mathfrak{A}$  is an object of  $\text{Src } \Upsilon$ .
- 2°.  $\langle \text{id}_c \rangle = (\lambda x \in \mathfrak{A} : x \sqcap c)$  for every  $c \in \mathfrak{A}$  where  $\mathfrak{A}$  is an object of  $\text{Src } \Upsilon$

DEFINITION 2207. *System of sides* is a system of presides which is both bounded and with identities.

<sup>1</sup>The idea for the name is that we consider one “side”  $\langle f \rangle$  of a funcooid instead of both sides  $\langle f \rangle$  and  $\langle f^{-1} \rangle$ .

PROPOSITION 2208.  $\langle 1_{\mathfrak{A}}^{\text{Src } \Upsilon} \rangle a = a$  for every system of presides.

PROOF. By properties of functors.  $\square$

DEFINITION 2209. I call a system of *monotone* presides a system of presides with additional axiom:

1°.  $\langle a \rangle$  is monotone.

DEFINITION 2210. I call a system of *distributive* presides a system of presides with additional axiom:

1°.  $\langle a \rangle (X \sqcup Y) = \langle a \rangle X \sqcup \langle a \rangle Y$ .

OBVIOUS 2211. Every distributive system of presides is monotone.

PROPOSITION 2212.  $\langle a \sqcap b \rangle X \sqsubseteq \langle a \rangle X \sqcap \langle b \rangle X$  for monotone systems of sides if Hom-sets are lattices.

DEFINITION 2213. A system of presides *with correct identities* is a system of presides with identities with additional axiom:

1°.  $\text{id}_b \circ \text{id}_a = \text{id}_{a \sqcap b}$ .

PROPOSITION 2214. Every faithful system of presides with identities is with correct identities.

PROOF.  $\langle \text{id}_b \circ \text{id}_a \rangle x = (\langle \text{id}_b \rangle \circ \langle \text{id}_a \rangle)x = \langle \text{id}_b \rangle \langle \text{id}_a \rangle x = b \sqcap a \sqcap x = \langle \text{id}_{b \sqcap a} \rangle x$ . Thus by faithfulness  $\text{id}_b \circ \text{id}_a = \text{id}_{b \sqcap a} = \text{id}_{a \sqcap b}$ .  $\square$

DEFINITION 2215. *Restricting* a side  $f$  to an object  $X$  is defined by the formula  $f|_X = f \circ \text{id}_X$ .

DEFINITION 2216. *Image* of a preside is defined by the formula  $\text{im } f = \langle f \rangle \top$ .

DEFINITION 2217. Protofunctors *over* a set  $X$  of functors is a protofunctor  $f$  such that  $\langle f \rangle \in X \wedge \langle f^{-1} \rangle \in X$ .

### 3. Concrete examples of sides

OBVIOUS 2218. The category  $\mathbf{Rel}$  with  $\langle f \rangle = \langle f \rangle^*$  for  $f \in \mathbf{Rel}$  and usual  $\text{id}_c$  defines a distributive system of sides with correct identities.

#### 3.1. Some subsides.

DEFINITION 2219. *Full subsystem* of a system  $\Upsilon$  of presides is the functor  $\Upsilon$  restricted to a full subcategory of  $\text{Src } \Upsilon$ .

OBVIOUS 2220. Full subsystem of a system of presides is always a system of presides.

OBVIOUS 2221. Full subsystem of a bounded system of presides is always a bounded subsystem of presides.

OBVIOUS 2222.

1°. Full subsystem of a system of presides with identities is always with identities.

2°. Full subsystem of a system of presides with correct identities is always with correct identities.

OBVIOUS 2223. Full subsystem of a distributive system of presides is always a distributive system of presides.

OBVIOUS 2224. Full subsystem of a system of sides is always a system of sides.

### 3.2. Functors and pointfree functors.

PROPOSITION 2225. The category of pointfree functors between starrish join-semilattices with usual  $\langle f \rangle$  defines a system of presides.

PROOF. Theorem 1632. □

PROPOSITION 2226. The category of pointfree functors between bounded starrish join-semilattices with usual  $\langle f \rangle$  defines a system of bounded presides.

PROOF. Take the proof of theorem 1629 into account. □

PROPOSITION 2227. The category of pointfree functors from a starrish join-semilattices to a separable starrish join-semilattices defines a distributive system of presides.

PROOF. Theorem 1604. □

PROPOSITION 2228. The category of pointfree functors between starrish lattices with usual  $\langle f \rangle$  and usual  $\text{id}_c$  defines a system of presides with correct identities.

PROOF. That it is with identities is obvious.

That it is with correct identities is obvious. □

OBVIOUS 2229. The category of pointfree functors between bounded starrish lattices with usual  $\langle f \rangle$  and usual  $\text{id}_c$  defines a system of sides with correct identities.

PROPOSITION 2230. The category of functors with usual  $\langle f \rangle$  and usual  $\text{id}_c$  defines a system of sides with correct identities.

PROOF. Because it can be considered a full subsystem of the category of point-free functors between bounded starrish lattices with usual  $\langle f \rangle$ . □

### 3.3. Galois connections.

PROPOSITION 2231. The category of Galois connections between (small) lattices with least elements together with usual  $\langle f \rangle$  defines a distributive system of presides.

PROOF. Propositions 2195 and 2196 for a system of presides.

It is distributive because lower adjoints preserve all joins. □

PROPOSITION 2232. The category of Galois connections between (small) bounded lattices together with usual  $\langle f \rangle$  defines a bounded system of presides.

PROOF. Theorem 2199. □

PROPOSITION 2233. The category of Galois connections between (small) Heyting lattices together with usual  $\langle f \rangle$  defines a system of sides with correct identities.

PROOF. Theorem 2200 ensures that they a system of sides with identities. The identities are correct due to faithfulness. □

### 3.4. Reloids.

PROPOSITION 2234. Reloids with the functor  $f \mapsto \langle (\text{FCD})f \rangle$  and usual  $\text{id}_c$  form a system of sides with correct identities.

PROOF. It is really a functor because  $\langle (\text{FCD})g \rangle \circ \langle (\text{FCD})f \rangle = \langle (\text{FCD})g \circ (\text{FCD})f \rangle = \langle (\text{FCD})(g \circ f) \rangle$  for every composable reloids  $f$  and  $g$ .

$$\langle a \rangle \perp = \langle (\text{FCD})a \rangle \perp = \perp;$$

$$\begin{aligned} \langle a \sqcup b \rangle X &= \langle (\text{FCD})(a \sqcup b) \rangle X = \langle (\text{FCD})a \sqcup (\text{FCD})b \rangle X = \\ & \langle (\text{FCD})a \rangle X \sqcup \langle (\text{FCD})b \rangle X = \langle a \rangle X \sqcup \langle b \rangle X; \end{aligned}$$



thus it is a system of presides.

That this is a bounded system of presides follows from the formulas  $(\text{FCD})_{\perp}^{\text{RLD}(A,B)} = \perp$  and  $(\text{FCD})_{\top}^{\text{RLD}(A,B)} = \top$ .

It is with identities, because proposition 1065. It is with correct identities by proposition 1025.  $\square$

**FiXme:** Also for pointfree reلودs.

**FiXme:** These examples works for (dagger) systems of sides with binary product.

#### 4. Product

DEFINITION 2235. *Binary product* of objects of presides with identities is defined by the formula  $X \times Y = \text{id}_Y \circ \top \circ \text{id}_X$ .

DEFINITION 2236. System of presides with identities is *with correct binary product* when  $f \sqcap (X \times Y) = \text{id}_Y \circ f \circ \text{id}_X$  for every preside  $f$ .

PROPOSITION 2237.  $\langle A \times B \rangle X = \begin{cases} \perp & \text{if } X \simeq A \\ B & \text{if } X \not\simeq A \end{cases}$

PROOF.

$$\begin{aligned} \langle A \times B \rangle X &= \langle \text{id}_B \circ \top \circ \text{id}_A \rangle X = \langle \text{id}_B \rangle \langle \top \rangle \langle \text{id}_A \rangle X = \\ &= B \sqcap \langle \top \rangle (X \sqcap A) = B \sqcap \begin{cases} \perp & \text{if } X \simeq A \\ \top & \text{if } X \not\simeq A \end{cases} = \begin{cases} \perp & \text{if } X \simeq A \\ B & \text{if } X \not\simeq A \end{cases} \end{aligned}$$

$\square$

DEFINITION 2238. I will call a system of sides *with correct meet* when

$$(X_0 \times Y_0) \sqcap (X_1 \times Y_1) = (X_0 \sqcap X_1) \times (Y_0 \sqcap Y_1).$$

PROPOSITION 2239. Faithful systems of presides with identities are with correct meet.

PROOF.  $(X_0 \times Y_0) \sqcap (X_1 \times Y_1) = \text{id}_{Y_1} \circ (X_0 \times Y_0) \circ \text{id}_{X_1}$ . Thus

$$\begin{aligned} \langle (X_0 \times Y_0) \sqcap (X_1 \times Y_1) \rangle P &= \langle \text{id}_{Y_1} \rangle \langle X_0 \times Y_0 \rangle \langle \text{id}_{X_1} \rangle P = \\ &= \langle \text{id}_{Y_1} \rangle \begin{cases} \perp & \text{if } X_0 \simeq \langle \text{id}_{X_1} \rangle P \\ Y_0 & \text{if } X_0 \not\simeq \langle \text{id}_{X_1} \rangle P \end{cases} = \begin{cases} \perp & \text{if } X_0 \sqcap X_1 \simeq P \\ Y_0 \sqcap Y_1 & \text{if } X_0 \sqcap X_1 \not\simeq P \end{cases} = \\ &= \langle (X_0 \sqcap X_1) \times (Y_0 \sqcap Y_1) \rangle P. \end{aligned}$$

So  $(X_0 \times Y_0) \sqcap (X_1 \times Y_1) = (X_0 \sqcap X_1) \times (Y_0 \sqcap Y_1)$  follows by full faithfulness.  $\square$

PROPOSITION 2240. Systems of presides with correct identities are with correct meet.

PROOF.  $(X_0 \times Y_0) \sqcap (X_1 \times Y_1) = \text{id}_{Y_1} \circ (X_0 \times Y_0) \circ \text{id}_{X_1} = \text{id}_{Y_1} \circ (\text{id}_{Y_0} \circ \top \circ \text{id}_{X_0}) \circ \text{id}_{X_1} = \text{id}_{Y_0 \sqcap Y_1} \circ \top \circ \text{id}_{X_0 \sqcap X_1} = (X_0 \sqcap X_1) \times (Y_0 \sqcap Y_1)$ .  $\square$

For some sides holds the formula  $f \circ (X \times Y) = X \times \langle f \rangle Y$ . I refrain to give a name for this property.

### 5. Negative results

The following negative result generalizes theorem 3.8 in [3].

**THEOREM 2241.** The element  $1^{(\text{Src } \Upsilon)(\mathfrak{A}, \mathfrak{A})}$  is not complemented if  $\mathfrak{A}$  is a non-atomic boolean lattice, for every monotone system of sides.

**PROOF.** Let  $T = 1^{(\text{Src } \Upsilon)(\mathfrak{A}, \mathfrak{A})}$ .

Let's suppose  $T \sqcup V = \top$  for  $V \in (\text{Src } \Upsilon)(\mathfrak{A}, \mathfrak{A})$  and prove  $T \sqcap V \neq \perp$ .

Then  $\langle T \sqcup V \rangle a = \top$  for all  $a \neq \perp$  and thus  $\langle V \rangle a \sqcup a = \top$ .

Consequently  $\langle V \rangle a \sqsupseteq \neg a$  for all  $a \neq \perp$ .

If  $a$  isn't an atom, then there exists  $b$  with  $0 \sqsubset b \sqsubset a$  and hence  $\langle V \rangle a \sqsupseteq \langle V \rangle b \sqsupseteq \neg b \sqsupseteq \neg a$ ; thus  $\langle V \rangle a \sqsupseteq \neg a$ .

There is such  $c \sqsubset \top$  that  $a \sqsubseteq c$  for every atom  $a$ . (Really, suppose some element  $p \neq \perp$  has no atoms. Thus all atoms are in  $\neg p$ .)

For  $a \not\sqsubseteq c$  we have  $\langle V \rangle a \sqcap a \sqsubset \perp$  for all  $a \sqsubseteq \neg c$  thus  $\langle T \sqcap V \rangle a \sqsupseteq \langle V \rangle a \sqcap a \sqsubset \perp$ .

Thus  $\langle (T \sqcap V) \circ \text{id}_{\neg c} \rangle a \sqsubset \perp$

So  $T \sqcap V \sqsupseteq (T \sqcap V) \circ \text{id}_{\neg c} \sqsubset \perp$ . So  $V$  is not a complement of  $T$ .  $\square$

**COROLLARY 2242.**  $(\text{Src } \Upsilon)(\mathfrak{A}, \mathfrak{A})$  is not boolean if  $\mathfrak{A}$  is a non-atomic boolean lattice.

### 6. Dagger systems of sides

**PROPOSITION 2243.**

- 1°. For a partially ordered dagger category, each of Hom-set of which has least element, we have  $\perp^\dagger = \perp$ .
- 2°. For a partially ordered dagger category, each of Hom-set of which has greatest element, we have  $\top^\dagger = \top$ .

**PROOF.**  $\forall f \in \text{Hom}(A, B) : \perp^\dagger \sqsubseteq f \Leftrightarrow \forall f \in \text{Hom}(A, B) : \perp \sqsubseteq f^\dagger \Leftrightarrow \forall f \in \text{Hom}(A, B) : \perp \sqsubseteq f \Leftrightarrow 1$ . Thus  $\perp^\dagger$  is the least.

The other items is dual.  $\square$

**DEFINITION 2244.** *Dagger system of presides with identities* is system of pre-sides with identities with category  $\text{Src } \Upsilon$  being a partially ordered dagger category and  $(\text{id}_X)^\dagger = \text{id}_X$  for every  $X$ .

**PROPOSITION 2245.** For a system of sides we have  $(X \times Y)^\dagger = Y \times X$ .

**PROOF.**  $(X \times Y)^\dagger = (\text{id}_Y \circ \top \circ \text{id}_X)^\dagger = \text{id}_X^\dagger \circ \top^\dagger \circ \text{id}_Y^\dagger = \text{id}_X \circ \top \circ \text{id}_Y = Y \times X$ .  $\square$

**FiXme:** Which properties of pointfree funcoids can be generalized for sides?

## Backward Functors

This is a preliminary partial draft.

Fix a family  $\mathfrak{A}$  of posets.

DEFINITION 2246. Let  $f$  be a staroid of filters  $\mathfrak{F}(\mathfrak{A}_i)$  on boolean lattices  $\mathfrak{A}_i$ . *Backward functor* for the argument  $k \in \text{dom } \mathfrak{A}$  of  $f$  is the functor  $\text{Back}(f, k)$  defined by the formula (for every  $X \in \mathfrak{A}_k$ )

$$\langle \text{Back}(f, k) \rangle X = \left\{ \frac{L \in \prod_{i \in \text{dom } \mathfrak{A}} \mathfrak{F}(\mathfrak{A}_i)}{X \in \langle f \rangle_k L} \right\}.$$

PROPOSITION 2247. Backward functor is properly defined.

PROOF.  $\langle \text{Back}(f, k) \rangle^*(X \sqcup Y) = \left\{ \frac{L \in \prod \mathfrak{A}}{X \sqcup Y \in \langle f \rangle_k L} \right\} = \left\{ \frac{L \in \prod \mathfrak{A}}{X \in \langle f \rangle_k L \vee Y \in \langle f \rangle_k L} \right\} = \left\{ \frac{L \in \prod \mathfrak{A}}{X \in \langle f \rangle_k L} \right\} \cup \left\{ \frac{L \in \prod \mathfrak{A}}{Y \in \langle f \rangle_k L} \right\} = \langle \text{Back}(f, k) \rangle^* X \cup \langle \text{Back}(f, k) \rangle^* Y. \quad \square$

OBVIOUS 2248. Backward functor is co-complete.

PROPOSITION 2249. If  $f$  is a principal staroid then  $\text{Back}(f, k)$  is a complete functor.

PROOF. ?? □

PROPOSITION 2250.  $f$  can be restored from  $\text{Back}(f, k)$  (for every fixed  $k$ ).

PROOF. ?? □

PROPOSITION 2251.  $f \mapsto \text{Back}(f, k)$  is an order isomorphism  $\text{Strd}^{\mathfrak{A}} \rightarrow \text{FCD}(\mathfrak{A}_k, \text{Strd}^{(\text{dom } \mathfrak{A}) \setminus \{k\}})$ .

PROOF. ?? □

## Quasi-atoms

DEFINITION 2252. *Quasi-atoms* functor  $\mathcal{A}$  is the functor  $A \rightarrow \text{atoms}^{\mathfrak{A}} A$  defined by the formula  $\langle \mathcal{A} \rangle^* X = \text{atoms}^{\mathfrak{A}} X$ .

This really defines a functor because  $\text{atoms}^{\mathfrak{A}} \perp = \emptyset$  and  $\text{atoms}^{\mathfrak{A}}(X \cup Y) = \text{atoms}^{\mathfrak{A}} X \cup \text{atoms}^{\mathfrak{A}} Y$ .

OBVIOUS 2253.  $\mathcal{A}$  is a co-complete functor.

PROPOSITION 2254.  $\langle \mathcal{A}^{-1} \rangle^* Y = \bigsqcup Y$ .

PROOF.  $Y \not\leq \langle \mathcal{A} \rangle^* X \Leftrightarrow Y \not\leq \text{atoms}^{\mathfrak{A}} X \Leftrightarrow \exists x \in \text{atoms}^{\mathfrak{A}} X, y \in Y : x \not\leq y \Leftrightarrow \exists y \in Y : X \not\leq y \Leftrightarrow$  (because  $X$  is a principal filter)  $\Leftrightarrow X \not\leq \bigsqcup Y$ .  $\square$

Note  $\langle \mathcal{A} \rangle^* \mathcal{X} = \prod_{X \in \text{up } \mathcal{X}} \text{atoms}^{\mathfrak{A}} X$ ;

$\langle \mathcal{A}^{-1} \rangle^* \mathcal{Y} = \prod_{Y \in \text{up } \mathcal{Y}} \bigsqcup Y$  ( $\mathcal{Y}$  is filter on the set of ultrafilters).

Can  $\text{atoms}^{\mathfrak{A}} \mathcal{X}$  be restored knowing  $\langle \mathcal{A} \rangle^* \mathcal{X}$ ? Can  $\bigsqcup \mathcal{Y}$  be restored knowing  $\langle \mathcal{A}^{-1} \rangle^* \mathcal{Y}$ ?

PROPOSITION 2255. (Provided that  $A$  is infinite)  $\mathcal{A}$  is not complete.

PROOF. Take a nonprincipal ultrafilter  $x$ . Then  $\langle \mathcal{A}^{-1} \rangle^* \{x\} = \bigsqcup \{x\} = x$  is a nonprincipal filter.  $\square$

CONJECTURE 2256. There is such filter  $\mathcal{X}$  that  $\langle \mathcal{A} \rangle^* \mathcal{X}$  is non-principal.

Does quasi-atoms functor define a more elegant replacement of  $\text{atoms}^{\mathfrak{A}}$ ? Does this concept have any use?

## Cauchy Filters on Reloids

In this chapter I consider *low filters* on reloids, generalizing Cauchy filters on uniform spaces. Using low filters, I define Cauchy-complete reloids, generalizing complete uniform spaces.

FiXme: I forgot to note that Cauchy spaces induce topological (or convergence) spaces.

### 1. Preface

Replace `\langle ... \rangle` with `\supfun{...}` in L<sup>A</sup>T<sub>E</sub>X.

This is a preliminary partial draft.

To understand this article you need first look into my book [2].

<http://math.stackexchange.com/questions/401989/>

[what-are-interesting-properties-of-totally-bounded-uniform-spaces](http://math.stackexchange.com/questions/401989/what-are-interesting-properties-of-totally-bounded-uniform-spaces)

[http://ncatlab.org/nlab/show/proximity+space#uniform\\_spaces](http://ncatlab.org/nlab/show/proximity+space#uniform_spaces) for a proof sketch that proximities correspond to totally bounded uniformities.

### 2. Low spaces

FiXme: Analyze <http://link.springer.com/article/10.1007/s10474-011-0136-9> (“A note on Cauchy spaces”), <http://link.springer.com/article/10.1007/BF00873992> (“Filter spaces”). It also contains references to some useful results, including (“On continuity structures and spaces of mappings” freely available at <https://eudml.org/doc/16128>) that the category FIL of filter spaces is isomorphic to the category of filter merotopic spaces (copy its definition).

DEFINITION 2257. A *lower set*<sup>1</sup> of filters on  $U$  (a set) is a set  $\mathcal{C}$  of filters on  $U$ , such that if  $\mathcal{G} \sqsubseteq \mathcal{F}$  and  $\mathcal{F} \in \mathcal{C}$  then  $\mathcal{G} \in \mathcal{C}$ .

REMARK 2258. Note that we are particularly interested in nonempty (= containing the improper filter) lower sets of filters. This does not match the traditional theory of Cauchy spaces (see below) which are traditionally defined as not containing empty set. Allowing them to contain empty set has some advantages:

- Meet of any lower filters is a lower filter.
- Some formulas become a little simpler.

DEFINITION 2259. I call *low space* a set together with a nonempty lower set of filters on this set. Elements of a (given) low space are called *Cauchy filters*.

DEFINITION 2260.  $\text{GR}(U, \mathcal{C}) = \mathcal{C}$ ;  $\text{Ob}(U, \mathcal{C}) = U$ .  $\text{GR}(U, \mathcal{C})$  is read as *graph of space*  $(U, \mathcal{C})$ . I denote  $\text{Low}(U)$  the set of graphs of low spaces on the set  $U$ . Similarly I will denote its subsets  $\text{ASJ}(U)$ ,  $\text{CASJ}(U)$ ,  $\text{Cau}(U)$ ,  $\text{CCau}(U)$  (see below).

FiXme: Should use “space structure” instead of “graph of space”, to match customary terminology.

<sup>1</sup>Remember that our orders on filters is the reverse to set theoretic inclusion. It could be called an *upper set* in other sources.

DEFINITION 2261. Introduce an order on graphs of low spaces and on low spaces:  $\mathcal{C} \sqsubseteq \mathcal{D} \Leftrightarrow \mathcal{C} \subseteq \mathcal{D}$  and  $(U, \mathcal{C}) \sqsubseteq (U, \mathcal{D}) \Leftrightarrow \mathcal{C} \subseteq \mathcal{D}$ .

OBVIOUS 2262. Every set of low spaces on some set is partially ordered.

### 3. Almost sub-join-semilattices

DEFINITION 2263. For a join-semilattice  $\mathfrak{A}$ , a *almost sub-join-semilattice* is such a set  $S \in \mathcal{P}\mathfrak{A}$ , that if  $\mathcal{F}, \mathcal{G} \in S$  and  $\mathcal{F} \not\sqsubseteq \mathcal{G}$  then  $\mathcal{F} \sqcup \mathcal{G} \in S$ .

DEFINITION 2264. For a complete lattice  $\mathfrak{A}$ , a *completely almost sub-join-semilattice* is such a set  $S \in \mathcal{P}\mathfrak{A}$ , that if  $\prod T \neq \perp^{\mathcal{F}(X)}$  then  $\prod T \in S$  for every  $T \in \mathcal{P}S$ .

OBVIOUS 2265. Every completely almost sub-join-semilattice is a almost sub-join-semilattice.

### 4. Cauchy spaces

DEFINITION 2266. A *reflexive* low space is a low space  $(U, \mathcal{C})$  such that  $\forall x \in U : \uparrow^U \{x\} \in \mathcal{C}$ .

DEFINITION 2267. The *identity* low space  $1^{\text{Low}(U)}$  on a set  $U$  is the low space with graph  $\left\{ \frac{\uparrow^U \{x\}}{x \in U} \right\}$ .

OBVIOUS 2268. A low space  $f$  is reflexive iff  $f \supseteq 1^{\text{Low}(\text{Ob } f)}$ .

DEFINITION 2269. An *almost sub-join space* is a low space whose graph is an almost sub-join-semilattice. I will denote such spaces as **ASJ**.

DEFINITION 2270. A *completely almost sub-join space* is a low space whose graph is a completely almost sub-join-semilattice. I will denote such spaces as **CASJ**.

DEFINITION 2271. A *precauchy space* (aka *filter space*) is a reflexive low space. I will denote such spaces as **preCau**.

DEFINITION 2272. A *Cauchy space* is a precauchy space which is also an almost sub-join space. I will denote such spaces as **Cau**.

DEFINITION 2273. A *completely Cauchy space* is a precauchy space which is also a completely almost sub-join space. I will denote such spaces as **CCau**.

OBVIOUS 2274. Every completely Cauchy space is a Cauchy space.

PROPOSITION 2275.  $a \sqcup \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\} b = a \sqcup b$  for  $a, b \in \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\}$ , provided that  $\mathcal{F}$  is a proper fixed Cauchy filter on an almost sub-join space.

PROOF.  $\mathcal{F}$  is proper. So we have  $a \sqcap b \supseteq \mathcal{F} \neq \perp^{\mathcal{F}(X)}$ . Thus  $a \sqcup b$  is a Cauchy filter and so  $a \sqcup b \in \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\}$ .  $\square$

PROPOSITION 2276.  $\prod \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\} S = \prod S$  for nonempty  $S \in \mathcal{P} \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\}$ , provided that  $\mathcal{F}$  is a proper fixed Cauchy filter on a completely almost sub-join space.

PROOF.  $\mathcal{F}$  is proper. So for every nonempty  $S \in \mathcal{P} \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\}$  we have  $\prod S \supseteq \mathcal{F} \neq \perp^{\mathcal{F}(X)}$ . Thus  $\prod S$  is a Cauchy filter and so  $\prod S \in \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \supseteq \mathcal{F}} \right\}$ .  $\square$

COROLLARY 2277. Every proper Cauchy filter is contained in a unique maximal Cauchy filter (for completely almost sub-join spaces).

PROOF. Let  $\mathcal{F}$  be a proper Cauchy filter. Then  $\bigsqcup \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \sqsupseteq \mathcal{F}} \right\} \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \sqsupseteq \mathcal{F}} \right\}$  (existing by the above proposition) is the maximal Cauchy filter containing  $\mathcal{F}$ .

Suppose another maximal Cauchy filter  $\mathcal{T}$  contains  $\mathcal{F}$ . Then  $\mathcal{T} \in \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \sqsupseteq \mathcal{F}} \right\}$  and thus  $\mathcal{T} = \bigsqcup \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \sqsupseteq \mathcal{F}} \right\} \left\{ \frac{\mathcal{X} \in \mathcal{C}}{\mathcal{X} \sqsupseteq \mathcal{F}} \right\}$ .  $\square$

## 5. Relationships with symmetric reloids

**FiXme:** Also consider relationships with funcoids.

DEFINITION 2278. Denote  $(\text{RLD})_{\text{Low}}(U, \mathcal{C}) = \bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in \mathcal{C}} \right\}$ .

DEFINITION 2279.  $(\text{Low})\nu$  (*low space* for endoreloid  $\nu$ ) is a low space on  $U$  such that

$$\text{GR}(\text{Low})\nu = \left\{ \frac{\mathcal{X} \in \mathcal{F}(U)}{\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq \nu} \right\}.$$

THEOREM 2280. If  $(U, \mathcal{C})$  is a low space, then  $(U, \mathcal{C}) = (\text{Low})(\text{RLD})_{\text{Low}}(U, \mathcal{C})$ .

PROOF. If  $\mathcal{X} \in \mathcal{C}$  then  $\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq (\text{RLD})_{\text{Low}}(U, \mathcal{C})$  and thus  $\mathcal{X} \in \text{GR}(\text{Low})(\text{RLD})_{\text{Low}}(U, \mathcal{C})$ . Thus  $(U, \mathcal{C}) \sqsubseteq (\text{Low})(\text{RLD})_{\text{Low}}(U, \mathcal{C})$ .

Let's prove  $(U, \mathcal{C}) \sqsupseteq (\text{Low})(\text{RLD})_{\text{Low}}(U, \mathcal{C})$ .

Let  $\mathcal{A} \in \text{GR}(\text{Low})(\text{RLD})_{\text{Low}}(U, \mathcal{C})$ . We need to prove  $\mathcal{A} \in \mathcal{C}$ .

Really  $\mathcal{A} \times^{\text{RLD}} \mathcal{A} \sqsubseteq (\text{RLD})_{\text{Low}}(U, \mathcal{C})$ . It is enough to prove that  $\exists \mathcal{X} \in \mathcal{C} : \mathcal{A} \sqsubseteq \mathcal{X}$ .

Suppose  $\nexists \mathcal{X} \in \mathcal{C} : \mathcal{A} \sqsubseteq \mathcal{X}$ .

For every  $\mathcal{X} \in \mathcal{C}$  obtain  $X_{\mathcal{X}} \in \mathcal{X}$  such that  $X_{\mathcal{X}} \notin \mathcal{A}$  (if for all  $X \in \mathcal{X}$  we have  $X_{\mathcal{X}} \in \mathcal{A}$ , then  $\mathcal{X} \sqsupseteq \mathcal{A}$  what is contrary to our supposition).

It is now enough to prove  $\mathcal{A} \times^{\text{RLD}} \mathcal{A} \not\sqsubseteq \bigsqcup \left\{ \frac{\uparrow^U X_{\mathcal{X}} \times^{\text{RLD}} \uparrow^U X_{\mathcal{X}}}{\mathcal{X} \in \mathcal{C}} \right\}$ .

Really,  $\bigsqcup \left\{ \frac{\uparrow^U X_{\mathcal{X}} \times^{\text{RLD}} \uparrow^U X_{\mathcal{X}}}{\mathcal{X} \in \mathcal{C}} \right\} = \uparrow^{\text{RLD}(U, U)} \bigcup \left\{ \frac{\uparrow^U X_{\mathcal{X}} \times^{\text{RLD}} \uparrow^U X_{\mathcal{X}}}{\mathcal{X} \in \mathcal{C}} \right\}$ . So our claim takes the form  $\bigcup \left\{ \frac{\uparrow^U X_{\mathcal{X}} \times^{\text{RLD}} \uparrow^U X_{\mathcal{X}}}{\mathcal{X} \in \mathcal{C}} \right\} \not\sqsubseteq \text{GR}(\mathcal{A} \times^{\text{RLD}} \mathcal{A})$  that is  $\forall A \in \mathcal{A} : \bigcup \left\{ \frac{\uparrow^U X_{\mathcal{X}} \times^{\text{RLD}} \uparrow^U X_{\mathcal{X}}}{\mathcal{X} \in \mathcal{C}} \right\} \not\sqsupseteq A \times A$  what is true because  $X_{\mathcal{X}} \not\sqsupseteq A$  for every  $A \in \mathcal{A}$ .  $\square$

REMARK 2281. The last theorem does not hold with  $\mathcal{X} \times^{\text{FCD}} \mathcal{X}$  instead of  $\mathcal{X} \times^{\text{RLD}} \mathcal{X}$  (take  $\mathcal{C} = \left\{ \frac{\{x\}}{x \in U} \right\}$  for an infinite set  $U$  as a counter-example).

REMARK 2282. Not every symmetric reloid is in the form  $(\text{RLD})_{\text{Low}}(U, \mathcal{C})$  for some Cauchy space  $(U, \mathcal{C})$ . The same Cauchy space can be induced by different uniform spaces. See <http://math.stackexchange.com/questions/702182/different-uniform-spaces-having-the-same-set-of-cauchy-filters>

PROPOSITION 2283.

1°.  $(\text{Low})f$  is reflexive iff endoreloid  $f$  is reflexive.

2°.  $(\text{RLD})_{\text{Low}}f$  is reflexive iff low space  $f$  is reflexive.

PROOF.

1°.  $f$  is reflexive  $\Leftrightarrow 1^{\text{RLD}} \sqsubseteq f \Leftrightarrow \forall x \in \text{Ob } f : \uparrow(\{x\} \times \{x\}) \sqsubseteq f \Leftrightarrow \forall x \in \text{Ob } f : \uparrow\{x\} \times^{\text{RLD}} \uparrow\{x\} \sqsubseteq f \Leftrightarrow \forall x \in \text{Ob } f : \uparrow\{x\} \in (\text{Low})f \Leftrightarrow (\text{Low})f$  is reflexive.

2°. Let  $f$  is reflexive. Then  $\forall x \in \text{Ob } f : \uparrow\{x\} \in f; \forall x \in \text{Ob } f : \uparrow\{x\} \times^{\text{RLD}} \uparrow\{x\} \sqsubseteq (\text{RLD})_{\text{Low}}f; \forall x \in \text{Ob } f : \uparrow(\{x\} \times \{x\}) \sqsubseteq (\text{RLD})_{\text{Low}}f; 1^{\text{RLD}} \sqsubseteq (\text{RLD})_{\text{Low}}f$ .

Let now  $(\text{RLD})_{\text{Low}}f$  be reflexive. Then  $f = (\text{Low})(\text{RLD})_{\text{Low}}f$  is reflexive.  $\square$

DEFINITION 2284. A *transitive* low space is such low space  $f$  that  $(\text{RLD})_{\text{Low}} f \circ (\text{RLD})_{\text{Low}} f = (\text{RLD})_{\text{Low}} f$ .

REMARK 2285. The composition  $(\text{RLD})_{\text{Low}} f \circ (\text{RLD})_{\text{Low}} f$  may be inequal to  $(\text{RLD})_{\text{Low}} \mu$  for all low spaces  $\mu$  (exercise!). Thus usefulness of the concept of transitive low spaces is questionable.

REMARK 2286. Every low space is “symmetric” in the sense that it corresponds to a symmetric reloid.

THEOREM 2287.  $(\text{Low})$  is the upper adjoint of  $(\text{RLD})_{\text{Low}}$ .

PROOF. We will prove  $(\text{Low})(\text{RLD})_{\text{Low}} f \sqsupseteq f$  and  $(\text{RLD})_{\text{Low}}(\text{Low})g \sqsubseteq g$  (that  $(\text{Low})$  and  $(\text{RLD})_{\text{Low}}$  are monotone is obvious).

Really:

$$\begin{aligned} \text{GR}(\text{Low})(\text{RLD})_{\text{Low}} f &= \text{GR}(\text{Low}) \bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in \text{GR } f} \right\} = \\ &= \left\{ \frac{\mathcal{Y} \in \mathcal{F} \text{ Ob}(f)}{\mathcal{Y} \times^{\text{RLD}} \mathcal{Y} \sqsubseteq \bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in \text{GR } f} \right\}} \right\} \supseteq \left\{ \frac{\mathcal{Y} \in \mathcal{F} \text{ Ob}(f)}{\mathcal{Y} \in \text{GR } f} \right\} = \text{GR } f; \\ (\text{RLD})_{\text{Low}}(\text{Low})g &= \bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in \text{GR}(\text{Low})g} \right\} = \bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in \mathcal{F}(\text{Ob } g), \mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq g} \right\} \sqsubseteq g. \quad \square \end{aligned}$$

COROLLARY 2288.

- 1°.  $(\text{RLD})_{\text{Low}} \bigsqcup S = \bigsqcup \langle (\text{RLD})_{\text{Low}} \rangle^* S$ ;
- 2°.  $(\text{Low}) \bigsqcap S = \bigsqcap \langle (\text{Low}) \rangle^* S$ .

Below it's proved that  $(\text{Low})$  and  $(\text{RLD})_{\text{Low}}$  can be restricted to completely almost sub-join spaces and symmetrically transitive reloids. Thus they preserve joins of (completely) almost sub-join spaces and meets of symmetrically transitive reloids. **FixMe: Check. FixMe: Move it to be below the definition.**

## 6. Lattices of low spaces

PROPOSITION 2289.  $\mu \sqsubseteq \nu \Leftrightarrow \forall \mathcal{X} \in \text{GR } \mu \exists \mathcal{Y} \in \text{GR } \nu : \mathcal{X} \sqsubseteq \mathcal{Y}$  for low filter spaces (on the same set  $U$ ).

PROOF.

$\Rightarrow$ .  $\mu \sqsubseteq \nu \Leftrightarrow \text{GR } \mu \subseteq \text{GR } \nu \Rightarrow \forall \mathcal{X} \in \text{GR } \mu \exists \mathcal{Y} \in \text{GR } \nu : \mathcal{X} = \mathcal{Y} \Rightarrow \forall \mathcal{X} \in \text{GR } \mu \exists \mathcal{Y} \in \text{GR } \nu : \mathcal{X} \sqsubseteq \mathcal{Y}$ .

$\Leftarrow$ . Let  $\forall \mathcal{X} \in \text{GR } \mu \exists \mathcal{Y} \in \text{GR } \nu : \mathcal{X} \sqsubseteq \mathcal{Y}$ . Take  $\mathcal{X} \in \text{GR } \mu$ . Then  $\exists \mathcal{Y} \in \text{GR } \nu : \mathcal{X} \sqsubseteq \mathcal{Y}$ . Thus  $\mathcal{X} \in \text{GR } \nu$ . So  $\text{GR } \mu \subseteq \text{GR } \nu$  that is  $\mu \sqsubseteq \nu$ . □

OBVIOUS 2290.

- 1°.  $(\text{RLD})_{\text{Low}}$  is an order embedding.
- 2°.  $(\text{Low})$  is an order homomorphism.

I will denote  $\bigsqcup, \bigsqcap, \sqcup, \sqcap$  the lattice operations on low spaces or graphs of low spaces.

PROPOSITION 2291.  $\bigsqcup S = \bigcup S$  for every set  $S$  of graphs of low spaces on some set.

PROOF. It's enough to prove that there is a low space  $\mu$  such that  $\text{GR } \mu = \bigcup S$ . In other words, it's enough to prove that  $\bigcup S$  is a nonempty lower set, but that's obvious. **FixMe: A little more detailed proof.** □



PROPOSITION 2292.  $\prod S = \left\{ \frac{\prod \text{im } P}{P \in \prod_{X \in S} X} \right\}$  for every set  $S$  of graphs of low spaces on some set.

PROOF. First prove that there is such low space  $\mu$  that  $\mu = \left\{ \frac{\prod \text{im } P}{P \in \prod_{X \in S} X} \right\}$ . In other words, we need to prove that  $\left\{ \frac{\prod \text{im } P}{P \in \prod_{X \in S} X} \right\}$  is a nonempty lower set. That it is nonempty is obvious. Let filter  $\mathcal{G} \subseteq \mathcal{F}$  and  $\mathcal{F} \in \left\{ \frac{\prod \text{im } P}{P \in \prod_{X \in S} X} \right\}$ . Then  $\mathcal{F} = \prod \text{im } P$  for a  $P \in \prod_{X \in S} X$  that is  $P(X) \in X$  for every  $X \in S$ . Take  $P' = (\mathcal{G} \cap) \circ P$ . Then  $P' \in \prod_{X \in S} X$  because  $P'(X) \in X$  for every  $X \in S$  and thus obviously  $\mathcal{G} = \prod \text{im } P'$  and thus  $\mathcal{G} \in \left\{ \frac{\prod \text{im } P}{P \in \prod_{X \in S} X} \right\}$ . So such  $\mu$  exists.

It remains to prove that  $\mu$  is the greatest lower bound of  $S$ .

$\mu$  is a lower bound of  $S$ . Really, let  $X \in S$  and  $Y \in X$ . Then exists  $P \in \prod_{X \in S} X$  such that  $P(X) = Y$  (taken into account that every  $X$  is nonempty) and thus  $\text{im } P \ni Y$  and so  $\prod \text{im } P \sqsubseteq Y$ , that is (proposition 2289)  $\mu \sqsubseteq X$ .

Let  $\nu$  be a lower bound of  $S$ . It remains to prove that  $\mu \sqsupseteq \nu$ , that is  $\forall Q \in \nu : Q = \prod \text{im } P$  for some  $P \in \prod_{X \in S} X$ . Take  $P = (\lambda X \in S : Q)$ . This  $P \in \prod_{X \in S} X$  because  $Q \in X$  for every  $X \in S$ .  $\square$

COROLLARY 2293.  $f \sqcap g = \left\{ \frac{F \sqcap G}{F \in f, G \in g} \right\}$  for every graphs  $f$  and  $g$  of low spaces (on some set).

### 6.1. Its subsets.

PROPOSITION 2294. The set of sub-join low spaces (on some fixed set) is meet-closed in the lattice of low spaces on a set.

PROOF. Let  $f, g$  be graphs of almost sub-join spaces (on some fixed set),  $f \sqcap g = \left\{ \frac{F \sqcap G}{F \in f, G \in g} \right\}$ .

If  $\mathcal{A}, \mathcal{B} \in f \sqcap g$  and  $\mathcal{A} \neq \mathcal{B}$ , then  $\mathcal{A}, \mathcal{B} \in f$  and  $\mathcal{A}, \mathcal{B} \in g$ . Thus  $\mathcal{A} \sqcup \mathcal{B} \in f$  and  $\mathcal{A} \sqcup \mathcal{B} \in g$  and so  $\mathcal{A} \sqcup \mathcal{B} \in f \sqcap g$ .  $\square$

COROLLARY 2295. The set of Cauchy spaces (on some fixed set), is meet-closed in the lattice of low spaces on a set.

PROPOSITION 2296. The set of completely almost sub-join spaces is meet-closed in the lattice of low spaces on a set.

PROOF. Let  $S$  be a set of graphs of almost completely sub-join low spaces (on some fixed set).  $\prod S = \left\{ \frac{\prod \text{im } P}{P \in \prod_{X \in S} X} \right\}$ .

If  $\mathcal{A}, \mathcal{B} \in \prod S$  and  $\mathcal{A} \neq \mathcal{B}$ , then  $\mathcal{A}, \mathcal{B} \in X$  for every  $X \in S$ . Thus  $\mathcal{A} \sqcup \mathcal{B} \in X$  and so  $\mathcal{A} \sqcup \mathcal{B} \in \prod S$ .  $\square$

COROLLARY 2297. The set of completely Cauchy spaces is meet-closed in the lattice of low spaces on a set.

From the above it follows:

OBVIOUS 2298. The following sets are complete lattices in our order:

- 1°. almost sub-join spaces, whose graphs are almost sub-join-semilattices;
- 2°. completely almost sub-join spaces;
- 3°. reflexive low spaces;
- 4°. precauchy spaces;

- 5°. Cauchy spaces;  
6°. completely Cauchy spaces.

Denote  $Z(f) = \left\{ \frac{F \sqcup G}{F \in f, G \in f, F \not\leq G} \right\} \cup \{\perp\}$  for every set  $f$  of filters (on some fixed set).

PROPOSITION 2299.  $Z(f) \supseteq f$  for every set  $f$  of filters.

PROOF. Consider for  $F \in f$  both cases  $F = \perp$  and  $F \neq \perp$ . □

LEMMA 2300. For graphs of low spaces  $f, g$  (on the same set)

$$Q = \bigcup S \cup Z\left(\bigcup S\right) \cup Z\left(Z\left(\bigcup S\right)\right) \cup \dots$$

is a graph of some almost sub-join space.

PROOF. That it is nonempty and a lower set of filters is obvious. It remains to prove that it is an almost sub-join-semilattice.

Let  $\mathcal{A}, \mathcal{B} \in Q$  and  $\mathcal{A} \not\leq \mathcal{B}$ . Then

$$\mathcal{A}, \mathcal{B} \in \underbrace{Z \dots Z}_{n \text{ times}}\left(\bigcup S\right)$$

for a natural  $n$ . Thus

$$\mathcal{A} \sqcup \mathcal{B} \in \underbrace{Z \dots Z}_{n+1 \text{ times}}\left(\bigcup S\right)$$

and so  $\mathcal{A} \sqcup \mathcal{B} \in Q$ . □

PROPOSITION 2301. Join on the lattice of graphs of almost sub-join spaces is described by the formula

$$\bigsqcup^{\text{ASJ}} S = \bigcup S \cup Z\left(\bigcup S\right) \cup Z\left(Z\left(\bigcup S\right)\right) \cup \dots$$

PROOF. The right part of the above formula  $\mu$  is a graph of an almost sub-join space (lemma).

That  $\mu$  is an upper bound of  $S$  is obvious.

It remains to prove that  $\mu$  is the least upper bound.

Suppose  $\nu$  is an upper bound of  $S$ . Then  $\nu \supseteq \bigcup S$ . Thus, because  $\nu$  is an almost sub-join-semilattice,  $Z(\nu) \subseteq \nu$ , likewise  $Z(Z(\nu)) \subseteq \nu$ , etc. Consequently  $Z(\bigcup S) \subseteq \nu$ ,  $Z(Z(\bigcup S)) \subseteq \nu$ , etc. So we have  $\mu \subseteq \nu$ . □

PROPOSITION 2302. FiXme: Should be merged with the previous proposition.

$$\bigsqcup^{\text{ASJ}} S = \left\{ \frac{F_0 \sqcup \dots \sqcup F_{n-1}}{F_0, \dots, F_{n-1} \in \bigcup S, F_0 \not\leq F_1 \wedge F_1 \not\leq F_2 \wedge \dots \wedge F_{n-2} \not\leq F_{n-1} \text{ for } n \in \mathbb{N}} \right\}.$$

REMARK 2303. We take  $F_0 \sqcup \dots \sqcup F_{n-1} = \perp$  for  $n = 0$ .

PROOF. Denote the right part of the above formula as  $R$ .

Suppose  $F \in R$ . Let's prove by induction that  $F \in Q$ . If  $F = \perp$  that's obvious. Suppose we know that  $F_0 \sqcup \dots \sqcup F_{n-1} \in Q$  that is for a natural  $m$

$$F_0 \sqcup \dots \sqcup F_{n-1} \in \underbrace{Z \dots Z}_{m \text{ times}}\left(\bigcup S\right)$$

for  $F_0, \dots, F_{n-1} \in \bigcup S$ ,  $F_0 \not\leq F_1 \wedge F_1 \not\leq F_2 \wedge \dots \wedge F_{n-2} \not\leq F_{n-1}$  and also  $F_{n-1} \not\leq F_n$ . Then  $F_0 \sqcup \dots \sqcup F_{n-1} \not\leq F_n$  and thus  $F_0 \sqcup \dots \sqcup F_{n-1} \sqcup F_n \in \underbrace{Z \dots Z}_{m+1 \text{ times}}\left(\bigcup S\right)$  that is

$F_0 \sqcup \dots \sqcup F_{n-1} \sqcup F_n \in Q$ . So  $F \in Q$  for every  $F \in R$ .

Now suppose  $F \in Q$  that is for a natural  $m$

$$F \in \underbrace{Z \dots Z}_{m \text{ times}} \left( \bigcup S \right).$$

Let's prove by induction that  $F = F_0 \sqcup \dots \sqcup F_{n-1}$  for some  $F_0, \dots, F_{n-1} \in \bigcup S$  such that  $F_0 \not\prec F_1 \wedge F_1 \not\prec F_2 \wedge \dots \wedge F_{n-2} \not\prec F_{n-1}$ . If  $m = 0$  then  $F \in \bigcup S$  and our promise is obvious. Let our statement holds for a natural  $m$ . Prove that it holds for

$$F' \in \underbrace{Z \dots Z}_{m+1 \text{ times}} \left( \bigcup S \right).$$

We have  $F' = Z(F)$  for some  $F = F_0 \sqcup \dots \sqcup F_{n-1}$  where  $F_0 \not\prec F_1 \wedge F_1 \not\prec F_2 \wedge \dots \wedge F_{n-2} \not\prec F_{n-1}$ . The case  $F' = \perp$  is easy. So we can assume  $F' = A \sqcup B$  where  $A, B \in F$  and  $A \not\prec B$ . By the statement of induction  $A = A_0 \sqcup \dots \sqcup A_{p-1}$ ,  $B = B_0 \sqcup \dots \sqcup B_{q-1}$  for natural  $p$  and  $q$ , where  $A_0 \not\prec A_1 \wedge A_1 \not\prec A_2 \wedge \dots \wedge A_{p-2} \not\prec A_{p-1}$ ,  $B_0 \not\prec B_1 \wedge B_1 \not\prec B_2 \wedge \dots \wedge B_{q-2} \not\prec B_{q-1}$ . Take  $j$  such that  $A \not\prec B_j$  and then take  $i$  such that  $A_i \not\prec B_j$ . Then (using symmetry of the relation  $\not\prec$ ) we have  $(A_0 \not\prec A_1 \wedge A_1 \not\prec A_2 \wedge \dots \wedge A_{p-2} \not\prec A_{p-1}) \wedge (A_{p-1} \not\prec A_{p-2} \not\prec \dots \wedge A_{i+1} \not\prec A_i) \wedge A_i \not\prec B_j \wedge (B_j \not\prec B_{j-1} \wedge \dots \wedge B_1 \not\prec B_0) \wedge (B_0 \not\prec B_1 \wedge B_1 \not\prec B_2 \wedge \dots \wedge B_{q-2} \not\prec B_{q-1})$ . So  $F' = A \sqcup B$  is representable as the join of a finite sequence of filters with each adjacent pair of filters in this sequence being intersecting. That is  $F' \in Q$ .  $\square$

PROPOSITION 2304. The lattice of Cauchy spaces (on some set) is a complete sublattice of the lattice of almost sub-join spaces.

PROOF. It's obvious, taking into account obvious 2268.  $\square$

$$\text{Denote } Z_\infty(f) = \left\{ \frac{\bigsqcup T}{T \in \mathcal{P}f \wedge \prod T \neq \perp} \right\} \cup \{\perp\}.$$

PROPOSITION 2305.  $Z_\infty(f) \supseteq f$ .

PROOF. Consider for  $F \in f$  both cases  $F = \perp$  and  $F \neq \perp$ .  $\square$

LEMMA 2306. If  $S$  is a set of graphs of low spaces, then

$$Q = \bigcup S \cup Z_\infty \left( \bigcup S \right) \cup Z_\infty \left( Z_\infty \left( \bigcup S \right) \right) \cup \dots$$

is a graph of a completely Cauchy space.

PROOF. That it is nonempty and a lower set of filters is obvious. It remains to prove that it is a completely almost sub-join-semilattice.

Let  $T \in \mathcal{P}Q$  and  $\prod T \neq \perp$ . Then

$$T \in \mathcal{P} \underbrace{Z_\infty \dots Z_\infty}_{n \text{ times}} \left( \bigcup S \right)$$

for a natural  $n$ . Thus

$$T \in \mathcal{P} \underbrace{Z_\infty \dots Z_\infty}_{n+1 \text{ times}} \left( \bigcup S \right)$$

and so  $\bigsqcup T \in Q$ .  $\square$

PROPOSITION 2307. The lattice of completely Cauchy spaces (on some set) is a complete sublattice of the lattice of completely almost sub-join spaces.

PROOF. It's obvious, taking into account obvious 2268.  $\square$

PROPOSITION 2308. Join of a set  $S$  on the lattice of graphs of completely almost sub-join-semilattice is described by the formula:

$$\text{CASJ} \quad \bigsqcup S = \bigcup S \cup Z_\infty \left( \bigcup S \right) \cup Z_\infty \left( Z_\infty \left( \bigcup S \right) \right) \cup \dots$$

PROOF. The right part of the above formula  $\mu$  is a graph of an almost sub-join space (lemma).

That  $\mu$  is an upper bound of  $S$  is obvious.

It remains to prove that  $\mu$  is the least upper bound.

Suppose  $\nu$  is an upper bound of  $S$ . Then  $\nu \supseteq \bigcup S$ . Thus, because  $\nu$  is an almost sub-join-semilattice,  $Z_\infty(\nu) \subseteq \nu$ , likewise  $Z_\infty(Z_\infty(\nu)) \subseteq \nu$ , etc. Consequently  $Z_\infty(\bigcup S) \subseteq \nu$ ,  $Z_\infty(Z_\infty(\bigcup S)) \subseteq \nu$ , etc. So we have  $\mu \sqsubseteq \nu$ .  $\square$

CONJECTURE 2309.

$$1^\circ. \bigsqcup^{\text{CASJ}} S = \left\{ \frac{\bigsqcup T_0 \sqcup \dots \sqcup T_{n-1}}{n \in \mathbb{N}, T_0, \dots, T_{n-1} \in \bigcup S,} \right\};$$

$$\left. \begin{array}{l} \prod T_0 \neq \perp \wedge \dots \wedge \prod T_{n-1} \neq \perp, \\ \bigsqcup T_0 \not\leq \bigsqcup T_1 \wedge \dots \wedge \bigsqcup T_{n-2} \not\leq \bigsqcup T_{n-1}. \end{array} \right\}$$

$$2^\circ. \bigsqcup^{\text{CASJ}} S = \left\{ \frac{\bigsqcup T_0 \sqcup \bigsqcup T_1 \sqcup \dots}{T_0, T_1, \dots \in \bigcup S,} \right\}$$

$$\left. \begin{array}{l} \prod T_0 \neq \perp \wedge \prod T_2 \neq \perp \wedge \dots, \bigsqcup T_0 \not\leq \bigsqcup T_1 \wedge \bigsqcup T_1 \not\leq \bigsqcup T_2 \wedge \dots \end{array} \right\}.$$

## 7. Up-complete low spaces

DEFINITION 2310. *Ideal base* is a nonempty subset  $S$  of a poset such that  $\forall a, b \in S \exists c \in S : (a, b \sqsubseteq c)$ .

OBVIOUS 2311. Ideal base is dual of filter base.

THEOREM 2312. Product of nonempty posets is a ideal base iff every factor is an ideal base.

PROOF. [FiXme: more detailed proof](#)

In one direction it is easy: Suppose one multiplier is not a dcpo. Take a chain with fixed elements (thanks our posets are nonempty) from other multipliers and for this multiplier take the values which form a chain without the join. This proves that the product is not a dcpo.

Let now every factor is dcpo.  $S$  is a filter base in  $\prod \mathfrak{A}$  iff each component is a filter base. Each component has a join. Thus by proposition 641  $S$  has a componentwise join.  $\square$

DEFINITION 2313. I call a low space *up-complete* when each ideal base (or equivalently every nonempty chain, see theorem 589) in this space has join in this space.

REMARK 2314. Elements of this ideal base are filters. (Thus is could be called a generalized ideal base.)

EXAMPLE 2315.

1 $^\circ$ .  $\left\{ \frac{\mathcal{X} \in \mathfrak{F}[0; +\infty[}{\exists \varepsilon > 0: \mathcal{X} \sqsubseteq \uparrow \varepsilon; +\infty[} \right\} \cup \uparrow \{0\}$  is a graph of Cauchy space on  $\mathbb{R}_+$ , but not up-complete.

2 $^\circ$ .  $\mathfrak{F}[0; +\infty[$  is a strictly greater graph of Cauchy space on  $\mathbb{R}_+$  and is up-complete.

LEMMA 2316. Let  $f$  be a reloid. Each ideal base  $T \subseteq \left\{ \frac{(A,B)}{\mathcal{A} \times^{\text{RLD}} \mathcal{B} \sqsubseteq f} \right\}$  has a join in this set.

PROOF. Let  $T$  be an ideal base and  $\forall (A, B) \in T : \mathcal{A} \times^{\text{FCD}} \mathcal{B} \sqsubseteq f$ .

$\forall (A, B) \in T \forall \mathcal{X} \in \mathcal{F} \text{ Src } f : (\mathcal{X} \not\prec \mathcal{A} \Rightarrow \mathcal{B} \sqsubseteq \langle f \rangle \mathcal{X})$ ;

taking join we have:

$\forall \mathcal{A} \in \text{Pr}_0 T \forall \mathcal{X} \in \mathcal{F} \text{ Src } f : (\mathcal{X} \not\prec \mathcal{A} \Rightarrow \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \sqsubseteq \langle f \rangle \mathcal{X})$ ;

$\forall \mathcal{A} \in \text{Pr}_0 T : \mathcal{A} \times^{\text{FCD}} \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \sqsubseteq f$ .

Now repeat a similar operation second time:

$\forall \mathcal{A} \in \text{Pr}_0 T : \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \times^{\text{FCD}} \mathcal{A} \sqsubseteq f^{-1}$ ;

$\forall \mathcal{A} \in \text{Pr}_0 T \forall \mathcal{Y} \in \mathcal{F} \text{ Dst } f : (\mathcal{Y} \not\prec \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \Rightarrow \mathcal{A} \sqsubseteq \langle f^{-1} \rangle \mathcal{Y})$ ;

$\forall \mathcal{Y} \in \mathcal{F} \text{ Dst } f : (\mathcal{Y} \not\prec \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \Rightarrow \bigsqcup_{\mathcal{A} \in \text{Pr}_0 T} \mathcal{A} \sqsubseteq \langle f^{-1} \rangle \mathcal{Y})$ ;

$\bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \times^{\text{FCD}} \bigsqcup_{\mathcal{A} \in \text{Pr}_0 T} \mathcal{A} \sqsubseteq f^{-1}$ ;

$\bigsqcup_{\mathcal{A} \in \text{Pr}_0 T} \mathcal{A} \times^{\text{FCD}} \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B} \sqsubseteq f$ . But  $\bigsqcup_{\mathcal{A} \in \text{Pr}_0 T} \mathcal{A} \times^{\text{FCD}} \bigsqcup_{\mathcal{B} \in \text{Pr}_1 T} \mathcal{B}$  is the join in consideration, because ideal base is ideal base in each argument.  $\square$

PROPOSITION 2317. A Cauchy space generated by an endoreloid is always up-complete.

PROOF. Let  $f$  be an endoreloid.  $\text{GR}(\text{Low})f = \left\{ \frac{\mathcal{X} \in \text{Ob } f}{\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq f} \right\}$ .

Let  $T \subseteq \left\{ \frac{\mathcal{X} \in \text{Ob } f}{\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq f} \right\}$  be an ideal base.

Then  $N = \left\{ \frac{(\mathcal{F}, \mathcal{F})}{\mathcal{F} \in T} \right\}$  is also an ideal base. Obviously  $N \subseteq \left\{ \frac{(A,B)}{\mathcal{A} \times^{\text{RLD}} \mathcal{B} \sqsubseteq f} \right\}$ . Thus by the lemma it has a join in  $\left\{ \frac{(A,B)}{\mathcal{A} \times^{\text{RLD}} \mathcal{B} \sqsubseteq f} \right\}$ . It's easy to see that this join is in  $\left\{ \frac{(A,A)}{\mathcal{A} \in \text{Ob } f, \mathcal{A} \times^{\text{RLD}} \mathcal{A} \sqsubseteq f} \right\}$ . Consequently  $T$  has a join in  $\left\{ \frac{\mathcal{X} \in \text{Ob } f}{\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq f} \right\}$ .  $\square$

It is long time known that (using our terminology) low space induced by a uniform space is a Cauchy space, but that it is complete and up-complete is probably first discovered by Victor Porton.

## 8. More on Cauchy filters

OBVIOUS 2318. Low filter on an endoreloid  $\nu$  is a filter  $\mathcal{F}$  such that

$$\forall U \in \text{GR } f \exists A \in \mathcal{F} : A \times A \subseteq U.$$

REMARK 2319. The above formula is the standard definition of Cauchy filters on uniform spaces.

PROPOSITION 2320. If  $\nu \sqsupseteq \nu \circ \nu^{-1}$  then every neighborhood filter is a Cauchy filter, that it

$$\nu \sqsupseteq \langle (\text{FCD})\nu \rangle^* \{x\} \times^{\text{RLD}} \langle (\text{FCD})\nu \rangle^* \{x\}$$

for every point  $x$ .

PROOF.  $\langle (\text{FCD})\nu \rangle^* \{x\} \times^{\text{RLD}} \langle (\text{FCD})\nu \rangle^* \{x\} = \langle (\text{FCD})\nu \rangle \uparrow^{\text{Ob } \nu} \{x\} \times^{\text{RLD}} \langle (\text{FCD})\nu \rangle \uparrow^{\text{Ob } \nu} \{x\} = \nu \circ (\uparrow^{\text{Ob } \nu} \{x\} \times^{\text{RLD}} \uparrow^{\text{Ob } \nu} \{x\}) \circ \nu^{-1} = \nu \circ (\uparrow^{\text{RLD}(\text{Ob } \nu, \text{Ob } \nu)} \{x, x\}) \circ \nu^{-1} \sqsubseteq \nu \circ \text{id}^{\text{RLD}(\text{Ob } \nu, \text{Ob } \nu)} \circ \nu^{-1} = \nu \circ \nu^{-1} \sqsubseteq \nu$ .  $\square$

PROPOSITION 2321. If  $\nu \sqsupseteq \nu \circ \nu^{-1}$  a filter converges (in  $\nu$ ) to a point, it is a low filter, provided that every neighborhood filter is a low filter.

PROOF. Let  $\mathcal{F} \sqsubseteq \langle (\text{FCD})\nu \rangle^* \{x\}$ . Then  $\mathcal{F} \times^{\text{RLD}} \mathcal{F} \sqsubseteq \langle (\text{FCD})\nu \rangle^* \{x\} \times^{\text{RLD}} \langle (\text{FCD})\nu \rangle^* \{x\} \sqsubseteq \nu$ .  $\square$

COROLLARY 2322. If a filter converges to a point, it is a low filter, provided that  $\nu \sqsupseteq \nu \circ \nu^{-1}$ .

### 9. Maximal Cauchy filters

LEMMA 2323. Let  $S$  be a set of sets with  $\prod \langle \uparrow^{\mathfrak{F}} \rangle^* S \neq 0^{\mathfrak{F}}$  (in other words,  $S$  has finite intersection property). Let  $T = \left\{ \frac{X \times X}{X \in S} \right\}$ . Then

$$\bigcup T \circ \bigcup T = \bigcup S \times \bigcup S.$$

PROOF. Let  $x \in \bigcup S$ . Then  $x \in X$  for some  $X \in S$ .  $\langle \bigcup T \rangle \{x\} \supseteq \uparrow X \supseteq \bigcap S \neq \emptyset$ . Thus

$$\langle \bigcup T \circ \bigcup T \rangle \{x\} = \langle \bigcup T \rangle \langle \bigcup T \rangle \{x\} \in \langle \uparrow^{\text{FCD}} \bigcup T \rangle \prod \langle \uparrow^{\mathfrak{F}} \rangle S \supseteq \bigsqcup \left\{ \frac{\langle \uparrow^{\text{FCD}}(X \times X) \rangle \prod \langle \uparrow^{\mathfrak{F}} \rangle S}{X \in S} \right\} = \bigsqcup \left\{ \frac{\uparrow^{\mathfrak{F}} X}{X \in S} \right\} = \bigsqcup \langle \uparrow^{\mathfrak{F}} \rangle S \text{ that is } \langle \bigcup T \circ \bigcup T \rangle \{x\} \supseteq \bigcup S. \quad \square$$

COROLLARY 2324. Let  $S$  be a set of filters (on some fixed set) with nonempty meet. Let

$$T = \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in S} \right\}$$

Then

$$\bigsqcup T \circ \bigsqcup T = \bigsqcup S \times^{\text{RLD}} \bigsqcup S.$$

$$\text{PROOF. } \bigsqcup T \circ \bigsqcup T = \prod \left\{ \frac{\uparrow^{\mathfrak{F}}(X \circ X)}{X \in \bigsqcup T} \right\}.$$

If  $X \in \bigsqcup T$  then  $X = \bigcup_{Q \in T} (P_Q \times P_Q)$  where  $P_Q \in Q$ . Therefore by the lemma we have

$$\bigcup \left\{ \frac{P_Q \times P_Q}{Q \in T} \right\} \circ \bigcup \left\{ \frac{P_Q \times P_Q}{Q \in T} \right\} = \bigcup_{Q \in T} P_Q \times \bigcup_{Q \in T} P_Q.$$

Thus  $X \circ X = \bigcup_{Q \in T} P_Q \times \bigcup_{Q \in T} P_Q$ .

$$\text{Consequently } \bigsqcup T \circ \bigsqcup T = \prod \left\{ \frac{\uparrow^{\mathfrak{F}}(\bigcup_{Q \in T} P_Q \times \bigcup_{Q \in T} P_Q)}{X \in \bigsqcup T} \right\} \supseteq \bigsqcup S \times^{\text{RLD}} \bigsqcup S.$$

$$\bigsqcup T \circ \bigsqcup T \sqsubseteq \bigsqcup S \times^{\text{RLD}} \bigsqcup S \text{ is obvious.} \quad \square$$

DEFINITION 2325. I call an endoreloid  $\nu$  *symmetrically transitive* iff for every symmetric endofunctor  $f \in \text{FCD}(\text{Ob } \nu, \text{Ob } \nu)$  we have  $f \sqsubseteq \nu \Rightarrow f \circ f \sqsubseteq \nu$ .

OBVIOUS 2326. It is symmetrically transitive if at least one of the following holds:

- 1°.  $\nu \circ \nu \sqsubseteq \nu$ ;
- 2°.  $\nu \circ \nu^{-1} \sqsubseteq \nu$ ;
- 3°.  $\nu^{-1} \circ \nu \sqsubseteq \nu$ .
- 4°.  $\nu^{-1} \circ \nu^{-1} \sqsubseteq \nu$ .

COROLLARY 2327. Every uniform space is symmetrically transitive.

PROPOSITION 2328.  $(\text{Low})\nu$  is a completely Cauchy space for every symmetrically transitive endoreloid  $\nu$ .

$$\text{PROOF. Suppose } S \in \mathcal{P} \left\{ \frac{\mathcal{X} \in \mathfrak{F}}{\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq \nu} \right\}.$$

$\bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in S} \right\} \sqsubseteq \nu$ ;  $\bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in S} \right\} \circ \bigsqcup \left\{ \frac{\mathcal{X} \times^{\text{RLD}} \mathcal{X}}{\mathcal{X} \in S} \right\} \sqsubseteq \nu$ ;  $\bigsqcup S \times^{\text{RLD}} \bigsqcup S \sqsubseteq \nu$  (taken into account that  $S$  has nonempty meet). Thus  $\bigsqcup S$  is Cauchy.  $\square$

PROPOSITION 2329. The neighbourhood filter  $\langle (\text{FCD})\nu \rangle^* \{x\}$  of a point  $x \in \text{Ob } \nu$  is a maximal Cauchy filter, if it is a Cauchy filter and  $\nu$  is a reflexive reloid.

**FiXme:** Does it holds for all low filters?

PROOF. Let  $\mathcal{N} = \langle (\text{FCD})\nu \rangle^* \{x\}$ . Let  $\mathcal{C} \sqsupseteq \mathcal{N}$  be a Cauchy filter. We need to show  $\mathcal{N} \sqsupseteq \mathcal{C}$ .

Since  $\mathcal{C}$  is Cauchy filter,  $\mathcal{C} \times^{\text{RLD}} \mathcal{C} \sqsubseteq \nu$ . Since  $\mathcal{C} \sqsupseteq \mathcal{N}$  we have  $\mathcal{C}$  is a neighborhood of  $x$  and thus  $\uparrow^{\text{Ob}\nu} \{x\} \sqsubseteq \mathcal{C}$  (reflexivity of  $\nu$ ). Thus  $\uparrow^{\text{Ob}\nu} \{x\} \times^{\text{RLD}} \mathcal{C} \sqsubseteq \mathcal{C} \times^{\text{RLD}} \mathcal{C}$  and hence  $\uparrow^{\text{Ob}\nu} \{x\} \times^{\text{RLD}} \mathcal{C} \sqsubseteq \nu$ ;

$$\mathcal{C} \sqsubseteq \text{im}(\nu|_{\uparrow^{\text{Ob}\nu} \{x\}}) = \langle (\text{FCD})\nu \rangle^* \{x\} = \mathcal{N}.$$

□

## 10. Cauchy continuous functions

DEFINITION 2330. A function  $f : U \rightarrow V$  is *Cauchy continuous* from a low space  $(U, \mathcal{C})$  to a low space  $(V, \mathcal{D})$  when  $\forall \mathcal{X} \in \mathcal{C} : \langle \uparrow^{\text{FCD}} f \rangle \mathcal{X} \in \mathcal{D}$ .

PROPOSITION 2331. Let  $f$  be a principal reloid. Then  $f \in \text{C}((\text{RLD})_{\text{Low}}\mathcal{C}, (\text{RLD})_{\text{Low}}\mathcal{D})$  iff  $f$  is Cauchy continuous.

$$\begin{aligned} f \circ (\text{RLD})_{\text{Low}}\mathcal{C} \circ f^{-1} &\sqsubseteq (\text{RLD})_{\text{Low}}\mathcal{D} \Leftrightarrow \\ \bigsqcup_{\mathcal{X} \in \mathcal{C}} (f \circ (\mathcal{X} \times^{\text{RLD}} \mathcal{X}) \circ f^{-1}) &\sqsubseteq (\text{RLD})_{\text{Low}}\mathcal{D} \Leftrightarrow \\ \bigsqcup_{\mathcal{X} \in \mathcal{C}} (\langle \uparrow^{\text{FCD}} f \rangle \mathcal{X} \times^{\text{RLD}} \langle \uparrow^{\text{FCD}} f \rangle \mathcal{X}) &\sqsubseteq (\text{RLD})_{\text{Low}}\mathcal{D} \Leftrightarrow \\ \forall \mathcal{X} \in \mathcal{C} : \langle \uparrow^{\text{FCD}} f \rangle \mathcal{X} \times^{\text{RLD}} \langle \uparrow^{\text{FCD}} f \rangle \mathcal{X} &\sqsubseteq (\text{RLD})_{\text{Low}}\mathcal{D} \Leftrightarrow \\ \forall \mathcal{X} \in \mathcal{C} : \langle \uparrow^{\text{FCD}} f \rangle \mathcal{X} &\in \mathcal{D}. \end{aligned}$$

Thus we have expressed Cauchy properties through the algebra of reloids.

## 11. Cauchy-complete reloids

DEFINITION 2332. An endoreloid  $\nu$  is *Cauchy-complete* iff every low filter for this reloid converges to a point.

REMARK 2333. In my book [2] *complete reloid* means something different. I will always prepend the word ‘‘Cauchy’’ to the word ‘‘complete’’ when meaning is by the last definition.

[https://en.wikipedia.org/wiki/Complete\\_uniform\\_space#Completeness](https://en.wikipedia.org/wiki/Complete_uniform_space#Completeness)

## 12. Totally bounded

<http://ncatlab.org/nlab/show/Cauchy+space>

DEFINITION 2334. Low space is called *totally bounded* when every proper filter contains a proper Cauchy filter.

OBVIOUS 2335. A reloid  $\nu$  is totally bounded iff

$$\forall X \in \mathcal{D} \text{ Ob}\nu \exists \mathcal{X} \in \mathfrak{F}^{\text{Ob}\nu} : (\perp \neq \mathcal{X} \sqsubseteq \uparrow^{\text{Ob}\nu} X \wedge \mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq \nu).$$

THEOREM 2336. A symmetric transitive reloid is totally bounded iff its Cauchy space is totally bounded.

PROOF.

$\Rightarrow$ . Let  $\mathcal{F}$  be a proper filter on  $\text{Ob}\nu$  and let  $a \in \text{atoms}\mathcal{F}$ . It’s enough to prove that  $a$  is Cauchy.

Let  $D \in \text{GR}\nu$ . Let also  $E \in \text{GR}\nu$  is symmetric and  $E \circ E \subseteq D$ . There exists a finite subset  $F \subseteq \text{Ob}\nu$  such that  $\langle E \rangle F = \text{Ob}\nu$ . Then obviously exists  $x \in F$  such that  $a \sqsubseteq \uparrow^{\text{Ob}\nu} \langle E \rangle \{x\}$ , but  $\langle E \rangle \{x\} \times \langle E \rangle \{x\} = E^{-1} \circ (\{x\} \times \{x\}) \circ E \subseteq D$ , thus  $a \times^{\text{RLD}} a \sqsubseteq \uparrow^{\text{RLD}(\text{Ob}\nu, \text{Ob}\nu)} D$ .

Because  $D$  was taken arbitrary, we have  $a \times^{\text{RLD}} a \sqsubseteq \nu$  that is  $a$  is Cauchy.

$\Leftarrow$ . Suppose that Cauchy space associated with a reloid  $\nu$  is totally bounded but the reloid  $\nu$  isn't totally bounded. So there exists a  $D \in \text{GR } \nu$  such that  $(\text{Ob } \nu) \setminus \langle D \rangle F \neq \emptyset$  for every finite set  $F$ .

Consider the filter base

$$S = \left\{ \frac{(\text{Ob } \nu) \setminus \langle D \rangle F}{F \in \mathcal{P} \text{Ob } \nu, F \text{ is finite}} \right\}$$

and the filter  $\mathcal{F} = \prod \langle \uparrow^{\text{Ob } \nu} \rangle S$  generated by this base. The filter  $\mathcal{F}$  is proper because intersection  $P \cap Q \in S$  for every  $P, Q \in S$  and  $\emptyset \notin S$ . Thus there exists a Cauchy (for our Cauchy space) filter  $\mathcal{X} \sqsubseteq \mathcal{F}$  that is  $\mathcal{X} \times^{\text{RLD}} \mathcal{X} \sqsubseteq \nu$ .

Thus there exists  $M \in \mathcal{X}$  such that  $M \times M \subseteq D$ . Let  $F$  be a finite subset of  $\text{Ob } \nu$ . Then  $(\text{Ob } \nu) \setminus \langle D \rangle F \in \mathcal{F} \sqsupseteq \mathcal{X}$ . Thus  $M \not\subseteq (\text{Ob } \nu) \setminus \langle D \rangle F$  and so there exists a point  $x \in M \cap ((\text{Ob } \nu) \setminus \langle D \rangle F)$ .

$\langle M \times M \rangle \{p\} \subseteq \langle D \rangle \{x\}$  for every  $p \in M$ ; thus  $M \subseteq \langle D \rangle \{x\}$ .

So  $M \subseteq \langle D \rangle (F \cup \{x\})$ . But this means that  $M \in \mathcal{X}$  does not intersect  $(\text{Ob } \nu) \setminus \langle D \rangle (F \cup \{x\}) \in \mathcal{F} \sqsupseteq \mathcal{X}$ , what is a contradiction (taken into account that  $\mathcal{X}$  is proper). □

<http://math.stackexchange.com/questions/104696/pre-compactness-total-boundedness-and-cauchy-sequential-compactness>

### 13. Totally bounded funcoids

DEFINITION 2337. A funcoid  $\nu$  is totally bounded iff

$$\forall X \in \text{Ob } \nu \exists \mathcal{X} \in \mathfrak{F}^{\text{Ob } \nu} : (0 \neq \mathcal{X} \sqsubseteq \uparrow^{\text{Ob } \nu} X \wedge \mathcal{X} \times^{\text{FCD}} \mathcal{X} \sqsubseteq \nu).$$

This can be rewritten in elementary terms (without using funcoidal product:  $\mathcal{X} \times^{\text{FCD}} \mathcal{X} \sqsubseteq \nu \Leftrightarrow \forall P \in \partial \mathcal{X} : \mathcal{X} \sqsubseteq \langle \nu \rangle P \Leftrightarrow \forall P \in \partial \mathcal{X}, Q \in \partial \mathcal{X} : P [\nu]^* Q \Leftrightarrow \forall P, Q \in \text{Ob } \nu : (\forall E \in \mathcal{X} : (E \cap P \neq \emptyset \wedge E \cap Q \neq \emptyset) \Rightarrow P [\nu]^* Q)$ .

Note that probably I am the first person which has written the above formula (for proximity spaces for instance) explicitly.

### 14. On principal low spaces

DEFINITION 2338. A low space  $(U, \mathcal{C})$  is *principal* when all filters in  $\mathcal{C}$  are principal.

PROPOSITION 2339. Having fixed a set  $U$ , principal reflexive low spaces on  $U$  bijectively correspond to principal reflexive symmetric endoreloids on  $U$ .

PROOF. ??

<http://math.stackexchange.com/questions/701684/union-of-cartesian-squares> □

### 15. Rest

[https://en.wikipedia.org/wiki/Cauchy\\_filter#Cauchy\\_filters](https://en.wikipedia.org/wiki/Cauchy_filter#Cauchy_filters)

[https://en.wikipedia.org/wiki/Uniform\\_space](https://en.wikipedia.org/wiki/Uniform_space) “Hausdorff completion of a uniform space” here)

<http://at.yorku.ca/z/a/a/b/13.htm> : the category **Prox** of proximity spaces and proximally continuous maps (i.e. maps preserving nearness between two sets) is isomorphic to the category of totally bounded uniform spaces (and uniformly continuous maps).



[https://en.wikipedia.org/wiki/Cauchy\\_space](https://en.wikipedia.org/wiki/Cauchy_space) <http://ncatlab.org/nlab/show/Cauchy+space>  
<http://arxiv.org/abs/1309.1748>  
[http://projecteuclid.org/download/pdf\\_1/euclid.pja/1195521991](http://projecteuclid.org/download/pdf_1/euclid.pja/1195521991)  
[http://www.emis.de/journals/HOA/IJMMS/Volume5\\_3/404620.pdf](http://www.emis.de/journals/HOA/IJMMS/Volume5_3/404620.pdf)  
[~/math/books/Cauchy\\_spaces.pdf](~/math/books/Cauchy_spaces.pdf)  
<https://ncatlab.org/nlab/show/Cauchy+space> defines compact Cauchy spaces!  
<http://www.hindawi.com/journals/ijmms/1982/404620/abs/> (open access article) describes criteria for a Cauchy space to be uniformizable.

## Funcoidal groups

REMARK 2340. **FiXme: Move this into the book.** If  $\mu$  and  $\nu$  are cocomplete endofunctors, then we can describe  $f \in C(\mu, \nu)$  without using filters by the formulas:

- 1°.  $\langle f \rangle^* \langle \mu \rangle^* X \sqsubseteq \langle \nu \rangle^* \langle f \rangle^* X$  (for every set  $X$  in  $\mathcal{P} \text{Ob } \mu$ )
- 2°.  $\langle \mu \rangle^* X \sqsubseteq \langle f^{-1} \rangle^* \langle \nu \rangle^* \langle f \rangle^* X$  (for every set  $X$  in  $\mathcal{P} \text{Ob } \mu$ )
- 3°.  $\langle f \rangle^* \langle \mu \rangle^* \langle f^{-1} \rangle^* Y \sqsubseteq \langle \nu \rangle^* Y$  (for every set  $Y$  in  $\mathcal{P} \text{Ob } \nu$ )

Funcoidal groups are modeled after topological groups (see Wikipedia) and are their generalization.

DEFINITION 2341. *Funcoidal group* is a group  $G$  together with endofunctor  $\mu$  on  $\text{Ob } G$  such that

- 1°.  $(y \cdot) \in C(\mu; \mu)$  for every  $y \in G$ ;
- 2°.  $(\cdot x) \in C(\mu; \mu)$  for every  $x \in G$ ;
- 3°.  $(x \mapsto x^{-1}) \in C(\mu; \mu)$  for every  $x \in G$ .

PROPOSITION 2342.  $t \mapsto y \cdot t \cdot x$  and  $t \mapsto y \cdot t^{-1} \cdot x$  are continuous functions.

PROOF. As composition of continuous functions. □

OBVIOUS 2343. Composition of functions of the forms  $t \mapsto y \cdot t \cdot x$  and  $t \mapsto y \cdot t^{-1} \cdot x$  are also a function of one of these forms.

What is the purpose of the following (yet unproved) proposition? I don't know, but it looks curious.

PROPOSITION 2344. Let  $E$  be a composition of functions of a form  $\langle \mu \rangle^*$ ,  $\langle y \cdot \rangle^*$ ,  $\langle \cdot x \rangle^*$ ,  $\langle^{-1} \rangle^*$  (where  $x$  and  $y$  vary arbitrarily) such that  $\mu$  is met in the composition at least once. Let also either  $\mu = \mu \circ \mu$  or  $\mu$  is met exactly once in the product. There are such elements  $x_0, y_0$  that either

- 1°.  $(t \mapsto y_0 \cdot t \cdot x_0) \circ \langle \mu \rangle \sqsubseteq E \sqsubseteq \langle \mu \rangle \circ (t \mapsto y_0 \cdot t \cdot x_0)$ ;
- 2°.  $(t \mapsto y_0 \cdot t^{-1} \cdot x_0) \circ \langle \mu \rangle \sqsubseteq E \sqsubseteq \langle \mu \rangle \circ (t \mapsto y_0 \cdot t^{-1} \cdot x_0)$ .

PROOF. Using continuity a few times we prove that  $E \sqsubseteq \langle \mu \rangle^* \circ \dots \circ \langle \mu \rangle^* \circ f_n \circ \dots \circ f_1$  where  $f_i$  are functions of the forms  $t \mapsto y \cdot t \cdot x$  or  $t \mapsto y \cdot t^{-1} \cdot x$  for  $n \in \mathbb{N}$ . But  $\langle \mu \rangle^* \circ \dots \circ \langle \mu \rangle^* = \langle \mu \rangle^*$  by conditions and  $f_n \circ \dots \circ f_1$  is of the form  $t \mapsto y \cdot t \cdot x$  or  $t \mapsto y \cdot t^{-1} \cdot x$  by above proposition.  $E \sqsubseteq \langle \mu \rangle \circ (t \mapsto y_0 \cdot t \cdot x_0)$  or  $E \sqsubseteq \langle \mu \rangle \circ (t \mapsto y_0 \cdot t^{-1} \cdot x_0)$

The second inequality is similar. Note that  $x_0$  and  $y_0$  are the same for the first and for the second item. □

$(G, \mu)$  vs  $(G, \mu^{-1})$  are they isomorphic?

**FiXme: We can also define reloidal groups.**

## 1. On “Each regular paratopological group is completely regular” article

In this chapter I attempt to rewrite the paper [1] in more general setting of functors and reloids. I attempt to construct a “royal road” to finding proofs of statements of this paper and similar ones, what is important because we lose 60 years waiting for any proof.

**1.1. Definition of normality.** By definition (slightly generalizing the special case if  $\mu$  is a quasi-uniform space from [1]) a pair of an endo-reloid  $\mu$  and a complete functor  $\nu$  (playing role of a generalization of a topological space) on a set  $U$  is *normal* when

$$\langle \nu^{-1} \rangle^* A \sqsubseteq \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle F \rangle^* A$$

for every entourage  $F \in \text{up } \mu$  of  $\mu$  and every set  $A \subseteq U$ .

Note that this is *not* the same as customary definition of normal topological spaces.

THEOREM 2345. An endoreloid  $\mu$  is normal on endoreloid  $\nu$  iff

$$\nu \circ \nu^{-1} \sqsubseteq \nu^{-1} \circ (\text{FCD})\mu.$$

PROOF. Equivalently transforming the criterion of normality (which should hold for all  $F \in \text{up } \mu$ ) using proposition 2189:

$$\langle \nu \rangle^* \langle \nu^{-1} \rangle^* A \sqsubseteq \langle \nu^{-1} \rangle^* \langle F \rangle^* A.$$

Also note

$$\prod_{F \in \text{up } \mu}^{\mathcal{F}} \langle \nu^{-1} \rangle^* \langle F \rangle^* A = (\text{because functors preserve filtered meets}) = \langle \nu^{-1} \rangle^* \prod_{F \in \text{up } \mu}^{\mathcal{F}} \langle F \rangle^* A = \langle \nu^{-1} \rangle^* \langle (\text{FCD})\mu \rangle^* A.$$

Thus the above is equivalent to  $\langle \nu \rangle^* \langle \nu^{-1} \rangle^* A \sqsubseteq \langle \nu^{-1} \rangle^* \langle (\text{FCD})\mu \rangle^* A$ .

And this is in turn equivalent to

$$\nu \circ \nu^{-1} \sqsubseteq \nu^{-1} \circ (\text{FCD})\mu.$$

□

DEFINITION 2346. An endofunctor  $\mu$  is *normal* on endofunctor  $\nu$  when  $\nu \circ \nu^{-1} \sqsubseteq \nu^{-1} \circ \mu$ . **FiXme: No need for  $\nu$  to be endomorphism.**

OBVIOUS 2347.

- 1°. Endoreloid  $\mu$  is normal on endofunctor  $\nu$  iff endofunctor  $(\text{FCD})\mu$  is normal on endofunctor  $\nu$ .
- 2°. Endofunctor  $\mu$  is normal on endoreloid  $\nu$  iff endofunctor  $(\text{RLD})_{\text{in}}\mu$  is normal on endofunctor  $\nu$ .

COROLLARY 2348. If  $\nu$  is a symmetric endofunctor and  $\mu \sqsupseteq \nu^{-1}$ , then it is normal.

COROLLARY 2349. (generalization of proposition 1 in [1]) If  $\nu$  is a symmetric endofunctor and  $\text{Compl } \mu \sqsupseteq \nu^{-1}$ , then it is normal.

DEFINITION 2350. A functor  $\nu$  is *normally reloidizable* iff there exist a reloid  $\mu$  such that  $(\mu, \nu)$  is normal and  $\nu = \text{Compl}(\text{FCD})\mu$ .

DEFINITION 2351. A functor  $\nu$  is *normally quasi-uniformizable* iff there exist a quasi-uniform space (= reflexive and transitive reloid)  $\mu$  such that  $(\mu, \nu)$  is normal and  $\nu = \text{Compl}(\text{FCD})\mu$ .

PROPOSITION 2352. A functor  $\nu$  is normally reloidizable iff there exist a functor  $\mu$  such that  $\mu$  is normal on  $\nu$  and  $\nu = \text{Compl } \mu$ .

PROPOSITION 2353. A funcoïd  $\nu$  is normally quasi-uniformizable iff there exist a quasi-proximity space (= reflexive and transitive funcoïd)  $\mu$  such that  $\mu$  is normal on  $\nu$  and  $\nu = \text{Compl } \mu$ .

PROOF. Obvious 2347 and the fact that (FCD) is an isomorphism between reflexive and transitive funcoïds and reflexive and transitive reloids.  $\square$

In other words, it is normally reloidazable or normally quasi-uniformizable when

$$(\text{Compl } \mu) \circ (\text{Compl } \mu)^{-1} \sqsubseteq (\text{Compl } \mu)^{-1} \circ \mu$$

for suitable  $\mu$ .

**1.2. Urysohn’s lemma and friends.** For a detailed proof of Urysohn’s lemma see also:

[http://homepage.math.uiowa.edu/~jsimon/COURSES/M132Fall07/UrysohnLemma\\_v5.pdf](http://homepage.math.uiowa.edu/~jsimon/COURSES/M132Fall07/UrysohnLemma_v5.pdf)

[https://proofwiki.org/wiki/Urysohn's\\_Lemma](https://proofwiki.org/wiki/Urysohn's_Lemma)

<http://planetmath.org/proofofurysohnslemma>

[https://en.wikipedia.org/wiki/Proximity\\_space](https://en.wikipedia.org/wiki/Proximity_space) says that “The resulting topology is always completely regular. This can be proven by imitating the usual proofs of Urysohn’s lemma, using the last property of proximal neighborhoods to create the infinite increasing chain used in proving the lemma.”

Below follows an alternative proof of Urysohn lemma. *The proof was based on a conjecture proved false, see example 1344!*

LEMMA 2354. If  $\langle \mu \rangle \mathcal{A} \asymp \mathcal{B}$  for a complete funcoïd  $\mu$  and  $\mathcal{A}, \mathcal{B}$  are filters on relevant sets, then there exists  $U \in \text{up } \mu$  such that  $\langle U \rangle \mathcal{A} \asymp \mathcal{B}$ .

PROOF. Prove that  $\left\{ \frac{\langle U \rangle \mathcal{A}}{U \in \text{up } \mu} \right\}$  is a filter base. That it is nonempty is obvious.

Let  $\mathcal{X}, \mathcal{Y} \in \left\{ \frac{\langle U \rangle \mathcal{A}}{U \in \text{up } \mu} \right\}$ . Then  $\mathcal{X} = \langle U_{\mathcal{X}} \rangle \mathcal{A}, \mathcal{Y} = \langle U_{\mathcal{Y}} \rangle \mathcal{A}$ . Because  $\mu$  is complete, we have (proposition 1124)  $U_{\mathcal{X}} \cap U_{\mathcal{Y}} \in \text{up } \mu$ . Thus  $\mathcal{X}, \mathcal{Y} \supseteq \langle U_{\mathcal{X}} \cap U_{\mathcal{Y}} \rangle \mathcal{A} \in \left\{ \frac{\langle U \rangle \mathcal{A}}{U \in \text{up } \mu} \right\}$ .

Thus  $\langle \mu \rangle \mathcal{A} \asymp \mathcal{B} \Leftrightarrow \mathcal{B} \cap \langle \mu \rangle \mathcal{A} = \perp \Leftrightarrow \exists U \in \text{up } \mu : \mathcal{B} \cap \langle U \rangle \mathcal{A} = \perp \Leftrightarrow \exists U \in \text{up } \mu : \langle U \rangle \mathcal{A} \asymp \mathcal{B}$ .  $\square$

COROLLARY 2355. If  $\langle \mu \rangle \mathcal{A} \asymp \langle \mu \rangle \mathcal{B}$  for a complete funcoïd  $\mu$  and  $\mathcal{A}, \mathcal{B}$  are filters on relevant sets, then there exists  $U \in \text{up } \mu$  such that  $\langle U \rangle \mathcal{A} \asymp \langle U \rangle \mathcal{B}$ .

PROOF. Applying the lemma twice we can obtain  $P, Q \in \text{up } \mu$  such that  $\langle P \rangle \mathcal{A} \asymp \langle Q \rangle \mathcal{B}$ . But because  $\mu$  is complete, we have  $U = P \cap Q \in \text{up } \mu$ , while obviously  $\langle U \rangle \mathcal{A} \asymp \langle U \rangle \mathcal{B}$ .  $\square$

LEMMA 2356. (assuming conjecture 1344) For every  $U \in \text{up } \mu$  (where  $\mu$  is a  $T_4$  topological space) such that  $\neg(A [U \circ U^{-1}]^* B)$  there is  $W \in \text{up } \mu$  such that  $U \circ U^{-1} \supseteq W \circ W^{-1} \circ W \circ W^{-1}$ . For it holds  $\neg(A [W \circ W^{-1}]^* B)$ . We can assume that  $\langle W \rangle^* X$  is open for every set  $X$ .

PROOF.  $U \circ U^{-1} \in \text{up}(\mu \circ \mu^{-1}) \subseteq \text{up}(\mu \circ \mu^{-1} \circ \mu \circ \mu^{-1})$  (normality used). Thus by the conjecture there exists  $W \in \text{up } \mu$  such that  $U \circ U^{-1} \supseteq W \circ W^{-1} \circ W \circ W^{-1}$ .  $W \circ W^{-1} \sqsubseteq U \circ U^{-1}$  thus  $\neg(A [W \circ W^{-1}]^* B)$ .

To prove that  $\langle W \rangle^* X$  is open for every set  $X$ , replace every  $\langle W \rangle^* \{x\}$  with an open neighborhood  $E \subseteq \langle W \rangle^* X$  of  $\langle \mu \rangle^* \{x\}$  (and note that union of open sets is open). This new  $W$  holds all necessary properties.  $\square$

LEMMA 2357. (assuming conjecture 1344) For every  $U \in \text{up } \mu$  (where  $\mu$  is a  $T_4$  topological space) such that  $\neg(A [U \circ U^{-1}]^* B)$  there is  $W \in \text{up } \mu$  such that  $U \circ U^{-1} \supseteq \mu^{-1} \circ W \circ W^{-1} \circ W \circ W^{-1}$ . For it holds  $\neg(A [W \circ W^{-1}]^* B)$ . We can assume that  $\langle W \rangle^* X$  is open for every set  $X$ .

PROOF. Applying the previous lemma twice, we have some open  $W \in \text{up } \mu$  such that

$$U \circ U^{-1} \supseteq W \circ W^{-1} \circ W \circ W^{-1} \circ W \circ W^{-1} \circ W \circ W^{-1}$$

and  $\neg(A [W \circ W^{-1}]^* B)$ . From this easily follows that

$$U \circ U^{-1} \supseteq \mu^{-1} \circ W \circ W^{-1} \circ W \circ W^{-1}.$$

□

A modified proof of Urysohn's lemma follows. This proof is in part based on [1]. (I attempt to find common generalization of Urysohn's lemma and results from [1]).

$$\mathbb{Q}_2 \stackrel{\text{def}}{=} \left\{ \frac{k/2^n}{k, n \in \mathbb{N}, 0 < k < 2^n} \right\}.$$

THEOREM 2358. Urysohn's lemma (see Wikipedia) for disjoint closed sets  $A$  and  $B$  and function  $f$  on a topological space  $\mu$  (considered as complete funcoid).

PROOF. (assuming conjecture 1344) (used ProofWiki among other sources)

Because  $A$  and  $B$  are disjoint closed sets, we have  $\langle \mu \rangle^* A \simeq \langle \mu \rangle^* B$ . Thus by the corollary 2355 take  $S_0 \in \text{up } \mu$  and  $\neg(A [S_0 \circ S_0^{-1}]^* B)$ .

We have  $\mu \circ \mu^{-1} \circ \mu \circ \mu^{-1} \subseteq \mu \circ \mu^{-1}$  that is  $\text{up}(\mu \circ \mu^{-1} \circ \mu \circ \mu^{-1}) \supseteq \text{up}(\mu \circ \mu^{-1})$ .

Let's prove by induction: There is a sequence  $S$  of binary relations starting with  $S_0$  such that  $\neg(A [S_i \circ S_i^{-1}]^* B)$  and  $S_i \circ S_i^{-1} \supseteq \mu^{-1} \circ S_{i+1} \circ S_{i+1}^{-1} \circ S_{i+1} \circ S_{i+1}^{-1}$ . It directly follows from the lemma (and uses the conjecture).

Denote  $U_i = S_{i+1} \circ S_{i+1}^{-1}$ . We have  $U_i \supseteq \mu^{-1} \circ U_{i+1} \circ U_{i+1}$  and  $\neg(A [U_i]^* B)$ .

By reflexivity of  $\mu$  we have  $U_{i+1} \subseteq U_{i+1} \circ U_{i+1} \subseteq U_i$ .

Define fractional degree of  $U$ :  $U^r \stackrel{\text{def}}{=} U_1^{r_1} \circ \dots \circ U_{l_r}^{r_{l_r}}$  for every  $r \in \mathbb{Q}_2$  where  $r_1 \dots r_{l_r}$  is the binary expansion of  $r$ .

Prove  $U_r \subseteq U_0$ . It is enough to prove  $U_0 \supseteq U_1 \circ \dots \circ U_{l_r}$ . It follows from  $U_2 \circ \dots \circ U_{l_r} \subseteq U_1, U_3 \circ \dots \circ U_{l_r} \subseteq U_2, \dots, U_{l_r} \subseteq U_{l_r-1}$  what was shown above.

Let's prove: For each  $p, q \in \mathbb{Q}_2$  such that  $p < q$  we have  $\mu^{-1} \circ U^p \subseteq U^q$ . We can assume binary expansion of  $p$  and  $q$  be the same length  $c$  (add zeros at the end of the shorter one). Now it is enough to prove

$$U_k \circ U_{k+1}^{q_{k+1}} \circ \dots \circ U_c^{q_c} \supseteq \mu^{-1} \circ U_{k+1}^{p_{k+1}} \circ U_{k+2}^{p_{k+2}} \circ \dots \circ U_c^{p_c}.$$

But for this it's enough

$$U_k \supseteq \mu^{-1} \circ U_{k+1} \circ U_{k+2} \circ \dots \circ U_c$$

what can be easily proved by induction: If  $k = c$  then it takes the form  $U_k \supseteq \mu^{-1}$  what is obvious. Suppose it holds for  $k$ . Then  $U_{k-1} \supseteq \mu^{-1} \circ U_k \circ U_k \supseteq \mu^{-1} \circ U_k \circ \mu^{-1} \circ U_{k+1} \circ U_{k+2} \circ \dots \circ U_c \supseteq \mu^{-1} \circ U_k \circ U_{k+1} \circ U_{k+2} \circ \dots \circ U_c$ , that is it holds for all natural  $k \leq c$ .

It is easy to prove that  $\langle U^r \rangle^* X$  is open for every set  $X$ .

We have  $\langle \mu^{-1} \rangle^* \langle U^p \rangle^* X \subseteq \langle U^q \rangle^* X$ .

$$f(z) \stackrel{\text{def}}{=} \inf \left( \{1\} \cup \left\{ \frac{q \in \mathbb{Q}_2}{z \in \langle U^q \rangle^* A} \right\} \right).$$

$f$  is properly defined because  $\{1\} \cup \left\{ \frac{q \in \mathbb{Q}_2}{z \in \langle U^q \rangle^* A} \right\}$  is nonempty and bounded.

If  $z \in A$  then  $z \in \langle U^q \rangle^* A$  for every  $q \in \mathbb{Q}_2$ , thus  $f(z) = 0$ , because obviously  $U^q \supseteq 1$ .

If  $z \in B$  then  $z \notin \langle U^q \rangle^* A$  for every  $q \in \mathbb{Q}_2$ , thus  $f(z) = 1$ , because  $U^q \subseteq U_0$ .

It remains to prove that  $f$  is continuous.

Let  $D(x) = \{1\} \cup \left\{ \frac{q \in \mathbb{Q}_2}{z \in \langle U^q \rangle^* A} \right\}$ .

To show that  $f$  is continuous, we first prove two smaller results:

(a)  $x \in \langle \mu^{-1} \rangle^* \langle U^r \rangle^* A \Rightarrow f(x) \leq r$ .

We have  $x \in \langle \mu^{-1} \rangle^* \langle U^r \rangle^* A \Rightarrow \forall s > r : x \in \langle U^s \rangle^* A$ , so  $D(x)$  contains all rationals greater than  $r$ . Thus  $f(x) \leq r$  by definition of  $f$ .

(b)  $x \notin \langle U^r \rangle^* A \Rightarrow f(x) \geq r$ .

We have  $x \notin \langle U^r \rangle^* A \Rightarrow \forall s < r : x \notin \langle U^s \rangle^* A$ . So  $D(x)$  contains no rational less than  $r$ . Thus  $f(x) \geq r$ .

Let  $x_0 \in S$  and let  $]c; d[$  be an open real interval containing  $f(x_0)$ . We will find a neighborhood  $T$  of  $x_0$  such that  $\langle f \rangle^* T \subseteq ]c; d[$ .

Choose  $p, q \in \mathbb{Q}$  such that  $c < p < f(x_0) < q < d$ . Let  $T = \langle U^q \rangle^* A \setminus \langle \mu^{-1} \rangle^* \langle U^p \rangle^* A$ .

Then since  $f(x_0) < q$ , we have that (b) implies vacuously that  $x \in \langle U^q \rangle^* A$ .

Since  $f(x_0) > p$ , (a) implies  $x_0 \notin \langle U^p \rangle^* A$ .

Hence  $x_0 \in T$ . Then  $T$  is a neighborhood of  $x_0$  because  $T$  is open.

Finally, let  $x \in T$ .

Then  $x \in \langle U^q \rangle^* A \subseteq \langle \mu^{-1} \rangle^* \langle U^q \rangle^* A$ . So  $f(x) \leq q$  by (a).

Also  $x \notin \langle \mu^{-1} \rangle^* \langle U^p \rangle^* A$ , so  $x \notin \langle U^p \rangle^* A$  and  $f(x) \geq p$  by (b).

Thus:  $f(x) \in [p; q] \subseteq ]c; d[$ .

Therefore  $f$  is continuous.

Claim A:  $f(x) > q \Rightarrow x \notin \langle \mu^{-1} \rangle^* \langle U^q \rangle^* A$

Claim B:  $f(x) < q \Rightarrow x \in \langle U^q \rangle^* A$

Proof of claim A: If  $f(x) > q$  then there must be some gap between  $q$  and  $D(x)$ ; in particular, there exists some  $q'$  such that  $q < q' < f(x)$ . But  $q' < f(x) \Rightarrow x \notin \langle U^{q'} \rangle^* A \Rightarrow x \notin \langle \mu^{-1} \rangle^* \langle U^{q'} \rangle^* A$  (using that  $\langle U^r \rangle^* X$  is open).

Proof of claim B: If  $f(x) < q$  then there exists  $q' \in D(x)$  such that  $f(x) < q' < q$ , in which case  $q \in D(x)$ , so  $x \in \langle U^q \rangle^* A$ .

To show that  $f$  is continuous, it's enough to prove that preimages of  $]a; 1]$  and  $[0; a[$  are open.

Suppose  $f(x) \in ]a; 1]$ . Pick some  $q$  with  $a < q < f(x)$ . We claim that the open set  $W = X \setminus \langle f^{-1} \rangle^* \langle U^q \rangle^* A$  is a neighborhood of  $x$  that is mapped by  $f$  into  $]a; 1]$ . First, by (A),  $f(x) > q \Rightarrow x \in W$ , so  $W$  is a neighborhood of  $x$ . If  $y$  is any point of  $W$ , then  $f(y)$  must be  $\geq q > a$ ; otherwise, if  $f(y) < q$ , then, by (B)  $y \in \langle U^q \rangle^* A \subseteq \langle f^{-1} \rangle^* \langle U^q \rangle^* A$ .

Suppose  $x \in f^{-1}[0; b[$  that is  $f(x) < b$  and pick  $q$  such that  $f(x) < q < b$ . By (B)  $x \in \langle U^q \rangle^* A$ . We claim that the neighborhood  $\langle U^q \rangle^* A$  is mapped by  $f$  into  $[0; b[$ . Suppose  $y$  is any point of  $\langle U^q \rangle^* A$ . Then  $q \in D(y)$ , so  $f(y) \leq q < b$ .  $\square$

**THEOREM 2359.** (from [1]) If  $\mu$  is a normal quasi-uniformity on a topological space  $\nu$ , then for any nonempty subset  $A \in \text{Ob } \nu$  and entourage  $U \in \text{up } \mu$  there exists a continuous function  $f : \text{Ob } \nu \rightarrow [0; 1]$  such that  $A \subseteq \langle f^{-1} \rangle^* \{0\} \subseteq \langle f^{-1} \rangle^* [0; 1] \subseteq \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U \rangle^* A$ .

**PROOF.** Choose inductively a sequence of entourages  $(U_n)_{n=0}^\infty$  such that  $U_0 = U$  and  $U_{n+1} \circ U_{n+1} \subseteq U_n$ .

Denote  $l_r = \max \left\{ \frac{n \in \mathbb{N}}{r_n = 1} \right\}$ .

Define  $U^r = U_{l_r}^{r_{l_r}} \circ \dots \circ U_1^{r_1}$

Prove  $\langle \nu^{-1} \rangle^* \langle U^q \rangle^* A \sqsubseteq \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U^r \rangle^* A$  for any  $q < r$  in  $\mathbb{Q}_2$ . **FixMe:** Can be easily rewritten with the formula  $\langle \nu \rangle^* \langle \nu^{-1} \rangle^* \langle U^q \rangle^* A \sqsubseteq \langle \nu^{-1} \rangle^* \langle U^r \rangle^* A$  instead. It may extend to non-complete functors.

There is such  $l$  that  $0 = q_l < r_l = 1$  and  $q_i = r_i$  for all  $i < l$ .

It follows  $l_q \neq l \leq l_r$ .

Consider variants:

$$\begin{aligned}
 l_q < l. \quad \langle \nu^{-1} \rangle^* \langle U^q \rangle^* A &\sqsubseteq \langle \nu^{-1} \rangle^* \langle U_{l_q} \circ \dots \circ U_1^{q_1 q_{l_q}} \rangle^* A = \\
 &\langle \nu^{-1} \rangle^* \langle U_{l_q}^{r_{l_q}} \circ \dots \circ U_1^{r_1} \rangle^* A \sqsubseteq \langle \nu^{-1} \rangle^* \langle U_{l-1}^{r_{l-1}} \circ \dots \circ U_1^{r_1} \rangle^* A \sqsubseteq \\
 &\langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_{l-1}^{r_{l-1}} \circ U_{l-1}^{r_{l-1}} \circ \dots \circ U_1^{r_1} \rangle^* A = \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U^r \rangle^* A \\
 &\text{(use } U_l^{r_l} \in \text{up(FCD)}\mu \text{ by theorem 992).} \\
 l < l_q. \text{ Inclusions } U_k \circ U_k \sqsubseteq U_{k-1} \text{ for } l < k \leq l_q + 1 \text{ guarantee that } U_{l_q+1} \circ U_{l_q} \circ \\
 &\dots \circ U_{l+1} \sqsubseteq U_l \text{ and then } \langle \nu^{-1} \rangle^* \langle U^q \rangle^* A \sqsubseteq \langle \nu^{-1} \rangle^* \langle U_{l_q}^{q_{l_q}} \circ \dots \circ U_1^{q_1} \rangle^* A \sqsubseteq \\
 &\langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_{l_q+1}^{q_{l_q+1}} \circ U_{l_q}^{q_{l_q}} \circ \dots \circ U_1^{q_1} \rangle^* A = \\
 &\langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_{l_q+1} \circ U_{l_q}^{q_{l_q}} \circ \dots \circ U_l^0 \circ \dots \circ U_1^{q_1} \rangle^* A \sqsubseteq \\
 &\langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_l \circ U_{l-1}^{q_{l-1}} \circ \dots \circ U_1^{q_1} \rangle^* A \sqsubseteq \\
 &\langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_l^{r_l} \circ U_{l-1}^{r_{l-1}} \circ \dots \circ U_1^{r_1} \rangle^* A \sqsubseteq \\
 &\langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_{l_r}^{r_{l_r}} \circ \dots \circ U_1^{r_1} \rangle^* A = \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U^r \rangle^* A.
 \end{aligned}$$

Define  $f$  by the formula  $f(z) = \inf \left( \{1\} \cup \left\{ \frac{q \in \mathbb{Q}_2}{z \in \langle \nu^{-1} \rangle^* \langle U^q \rangle^* A} \right\} \right)$ .

It is clear?? that  $A \sqsubseteq \langle f^{-1} \rangle^* \{0\}$  and  $\langle f^{-1} \rangle^* [0; 1[ \sqsubseteq \bigcup_{q \in \mathbb{Q}_2} \langle \nu^{-1} \rangle^* \langle U^q \rangle^* A = \bigcup_{r \in \mathbb{Q}_2} \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U^r \rangle^* A \sqsubseteq \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U_0 \rangle^* A$ .

To prove that the map  $f : X \rightarrow [0, 1]$  is continuous, it suffices to check that for every real number  $a \in ]0; 1[$  the sets  $\langle f^{-1} \rangle^* [0; a[$  and  $\langle f^{-1} \rangle^* ]a; 1]$  are open. This follows from the equalities

$$\langle f^{-1} \rangle^* [0; a[ = \bigcup_{\mathbb{Q}_2 \ni q < a} \langle \nu^{-1 \circ} \rangle^* \langle \nu^{-1} \rangle^* \langle U^q \rangle^* A \text{ and } \langle f^{-1} \rangle^* ]a; 1] = \bigcup_{\mathbb{Q}_2 \ni r > a} (X \setminus \langle \nu^{-1} \rangle^* \langle U^r \rangle^* A). \quad \square$$

How the formulas for normal ( $T_4$ ) topological spaces and normal quasi-uniformities are related? Maybe this works: Replacing  $\nu \rightarrow \mu \circ \mu^{-1}$ ,  $\mu \rightarrow 1$  makes  $\nu \circ \nu^{-1} \sqsubseteq \nu^{-1} \circ (\text{FCD})\mu \rightarrow \mu \circ \mu^{-1} \circ \mu \circ \mu^{-1} \sqsubseteq \mu \circ \mu^{-1}$ .

<https://www.researchgate.net/project/The-lattice-LG-of-group-topologies>

## Micronization

I defined “micronization” wrongly in my book and did some erroneous proofs about it. Here is an attempt to salvage it.

[https://en.wikipedia.org/wiki/Transitive\\_reduction](https://en.wikipedia.org/wiki/Transitive_reduction) is a special case of micronization. (Hm, maybe them coincide only for finite sets?)

DEFINITION 2360. *Micronization*  $\mu(E)$  of a binary relation  $E$  is defined by the formula:

$$\mu(E) = \prod^{\text{RLD}} \left\{ \frac{f \in \text{RLD}}{S^*(f) \supseteq E \wedge f \asymp f^2} \right\}$$

It’s wrong (consider micronization of  $\leq$  on real numbers (which should be addition of infinite small)).

QUESTION 2361. Under which conditions  $S^*(\mu(E)) = E$ ?



## More on connectedness

### 1. For topological spaces

PROPOSITION 2362. The following are pairwise equivalent:

- 1°. a topological space on a set  $U$  is connected. **FiXme: definition; can the topological definition be generalized to filters?**
- 2°.  $U$  is connected regarding  $f \sqcup f^{-1}$  if  $f$  is the corresponding complete functor.
- 3°.  $U$  is connected regarding  $f \sqcup f^{-1}$  if  $f$  is the corresponding closure space.
- 4°.  $U$  is connected regarding  $f \circ f^{-1}$  if  $f$  is the corresponding complete functor.

PROOF. ?? □

PROPOSITION 2363. There are filters  $\mathcal{A}, \mathcal{B}$ , such that there are no filters  $\mathcal{X} \sqsubseteq \mathcal{A}$ ,  $\mathcal{Y} \sqsubseteq \mathcal{B}$  such that  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A} \sqcup \mathcal{B}$  and  $\mathcal{X} \asymp \mathcal{Y}$ .

PROOF. <https://math.stackexchange.com/questions/2639206>

(It also follows that sometimes  $Z(Da)$  is not a complete lattice, because otherwise we could prove this theorem.) □

PROPOSITION 2364. If  $\mathcal{A}, \mathcal{B}$  are filters and  $\mathcal{A} \sqcup \mathcal{B} = U$  is principal filter, then there are sets  $X \sqsubseteq \mathcal{A}, Y \sqsubseteq \mathcal{B}$  such that  $X \sqcup Y = U$  and  $X \asymp Y$ .

PROOF. Take  $X = \text{Cor } \mathcal{A}$  and  $Y' = \text{Cor } \mathcal{B}$ . Then  $X \sqcup Y' = U$  because of co-separability of  $\mathfrak{F}(U)$ . Take  $Y = U \setminus X$ . Then  $X \sqcup Y = U$  and  $X \asymp Y$ . □

PROPOSITION 2365. A principal filter  $A$  is connected regarding endofunctor  $\mu$  iff

$$\forall X, Y \in \mathcal{P}(\text{Ob } \mu) \setminus \{\perp\} : (X \sqcup Y = A \wedge X \asymp Y \Rightarrow X [\mu] Y).$$

PROOF. Easily follows from ?? □

DEFINITION 2366. *Connected component* of a filter regarding a functor or a reloid is a maximal connected subfilter of this filter.

OBVIOUS 2367. Subfilter of a connected filter is connected.

PROPOSITION 2368. If  $U$  is a principal filter, then it is connected regarding  $\mu$  iff it is connected regarding  $S(\mu)$ . **FiXme: It should be presented as a corollary of a below theorem.**

PROOF. If  $U$  is connected regarding  $\mu$ , it is connected regarding  $S(\mu)$ , obviously.

Suppose  $U$  is connected regarding  $S(\mu)$ . Then for  $X, Y \in \mathcal{P}(\text{Ob } \mu) \setminus \{\perp\}$  we have if  $X \sqcup Y = U$  and  $X \asymp Y$ , then  $X [S(\mu)] Y$ . So  $X \times Y \neq 1 \sqcup \mu \sqcup \mu^2 \sqcup \dots$  and thus by distributivity for principal filter we have  $X \times Y \neq \mu^n$  for some  $n \geq ??$  that is  $X [\mu^n] Y$  for some  $n$  and thus there are atomic filters  $p_0, \dots, p_n$  such that  $p_0 \in \text{atoms}^{\mathfrak{S}} X$ ,  $p_n \in \text{atoms}^{\mathfrak{S}} Y$  and  $p_i [\mu] p_{i+1}$ . Thus there is  $k$  such that  $p_k [\mu] p_{k+1}$  and  $p_k \in \text{atoms}^{\mathfrak{S}} X$ ,  $p_{k+1} \in \text{atoms}^{\mathfrak{S}} Y$ . Thus  $X [\mu] Y$ . We have  $U$  connected regarding  $\mu$ . □

Also for  $S^*$

EXAMPLE 2369. Connected components may not form a weak partition.

PROOF. Consider funcoid  $1^{\text{FCD}(\mathbb{R})} \sqcup (\Delta \times^{\text{FCD}} \Delta)$  on real line. Then connected components are (prove!) non-zero singletons and  $\Delta$ . It is not a weak partition.  $\square$

CONJECTURE 2370. If the set of connected components is finite, then it is a strong partition. Moreover the set of connected components is a tearing.

Add more counter-examples (for non-principal filters).

OBVIOUS 2371. Improper filter  $\perp^{\mathcal{F}}$  is connected regarding:

- 1°. every funcoid;
- 2°. every reloid.

PROPOSITION 2372. The only filter connected regarding

- 1°.  $\perp^{\text{FCD}(A)}$ ;
- 2°.  $\perp^{\text{RLD}(A)}$

is the improper filter  $\perp^{\mathcal{F}}$ .

PROOF.

- 1°. Let  $\mathcal{A}$  be a filter. Take  $\mathcal{X} = \mathcal{Y} = \mathcal{A} \in \mathcal{F}(\text{Ob } \mu) \setminus \{\perp\}$ . Then  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A}$  but not  $\mathcal{X} [\mu] \mathcal{Y}$ .
- 2°.  $S_1^*(\perp^{\text{RLD}(A)}) = S_1(\perp^{\text{RLD}(A)}) = \perp^{\text{RLD}(A)}$ . Thus the only connected filter is  $\perp^{\mathcal{F}}$ .

$\square$

PROPOSITION 2373. Connected filters regarding

- 1°.  $1^{\text{FCD}(A)}$ ;
- 2°.  $1^{\text{RLD}(A)}$

are exactly ultrafilters and the improper filter.

PROOF. 1. That ultrafilters are connected follows from the fact that for every non-least  $\mathcal{X}, \mathcal{Y}$  such that  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A}$  we have  $\mathcal{X} = \mathcal{Y} = \mathcal{A}$  and thus  $\mathcal{X} [1^{\text{FCD}(A)}] \mathcal{Y}$ . So ultrafilters are connected; so is improper filter too, because improper filter is always connected.

It remains to prove that filters containing more than one distinct ultrafilter are not connected. Really let distinct ultrafilters  $a, b \in \text{atoms } \mathcal{A}$ . Then not  $a [1^{\text{FCD}(A)}] b$ . Thus  $\mathcal{A}$  is not connected.

2. A filter  $a$  is connected iff  $S_1^*(1^{\text{RLD}(A)} \sqcap (a \times^{\text{RLD}} a)) \sqsupseteq a \times^{\text{RLD}} a$  that is iff  $S_1^*(\text{id}_a^{\text{RLD}}) \sqsupseteq a \times^{\text{RLD}} a$ ,  
 $\prod_{F \in \text{up id}_a^{\text{RLD}}} S_1(F) \sqsupseteq a \times^{\text{RLD}} a$  what by properties of generalized filter bases is equivalent to  $\prod_{A \in \text{up } a} S_1(\text{id}_A) \sqsupseteq a \times^{\text{RLD}} a$ ;  $\prod_{A \in \text{up } a} \text{id}_A \sqsupseteq a \times^{\text{RLD}} a$ ;  $\text{id}_a^{\text{RLD}} \sqsupseteq a \times^{\text{RLD}} a$ . This is true exactly for ultrafilters and the improper filter.  $\square$

DEFINITION 2374. A *path* regarding funcoid  $\mu$  is a tuple  $p_0, \dots, p_n$  ( $n \in \mathbb{N}$ ) of atomic filters such that  $p_i [\mu] p_{i+1}$  for every  $i = 0, \dots, n-1$ .

The number  $n$  is called *path length*.

A path is *between* atomic filters  $a$  and  $b$  iff  $p_0 = a$  and  $p_n = b$ .

EXAMPLE 2375.  $\mu \sqsupseteq \text{id}_{\mathcal{A}}^{\text{FCD}}$  is not necessary for a filter  $\mathcal{A}$  to be connected regarding a funcoid  $\mu$ . Moreover  $\mu \sqsupseteq 1^{\text{FCD}}$  is not necessary for a filter  $\top$  to be connected regarding a funcoid  $\mu$ .

PROOF. For counterexample take  $\mu = \top \setminus 1$ .

$\langle \mu \rangle \{x\} = \top \setminus \{x\}$  (thus  $\mu \not\sqsupseteq 1^{\text{FCD}}$ ) and  $\langle \mu \rangle a = \top$  for a nontrivial ultrafilter  $a$ .

Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{F}(\text{Ob } \mu) \setminus \{\perp\}$  and  $\mathcal{X} \sqcup \mathcal{Y} = \top$ . If  $\mathcal{X}$  is a trivial ultrafilter then  $\langle \mu \rangle \mathcal{X} = \top \setminus \{x\}$  and thus  $\langle \mu \rangle \mathcal{X} \neq \mathcal{Y}$ , otherwise  $\langle \mu \rangle \mathcal{X} \neq \mathcal{Y}$ . So in any case  $\mathcal{X} [\mu] \mathcal{Y}$ . Funcoid  $\mu$  is connected.  $\square$

PROPOSITION 2376. If there is a nonzero-length path regarding  $\mu$  in the filter  $\mathcal{A}$  between any two its atomic subfilters, then it is connected regarding  $\mu$ .

PROOF. Let  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A}$ ,  $\mathcal{X} \neq \perp$ ,  $\mathcal{Y} \neq \perp$ . Let  $p_0, \dots, p_n$  ( $n \geq 1$ ) be a path in  $\mathcal{A}$  and  $p_0 \in \text{atoms } \mathcal{X}$  and  $p_n \in \text{atoms } \mathcal{Y}$ . Then (take  $k = \min\{i \in \{0, \dots, n-1\} \mid p_{i+1} \in \text{atoms } \mathcal{Y}\}$ ) there are  $p_k, p_{k+1}$  such that  $p_k \in \text{atoms } \mathcal{X}$ ,  $p_{k+1} \in \text{atoms } \mathcal{Y}$ . But  $p_k [\mu] p_{k+1}$  by definition of path. Thus  $\mathcal{X} [\mu] \mathcal{Y}$ .  $\square$

PROPOSITION 2377. If a filter  $\mathcal{A}$  is connected regarding funcoid  $\mu$  reflexive on  $\mathcal{A}$  then it is connected regarding every  $\mu^n$  for  $n \in \mathbb{Z}_+$ .

PROOF. Let  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A}$ ,  $\mathcal{X} \neq \perp$ ,  $\mathcal{Y} \neq \perp$ . We have  $\langle \mu \rangle \mathcal{X} \neq \mathcal{Y}$ .

Then  $\langle \mu \rangle \mathcal{X} \not\sqsupseteq \mathcal{X}$ ; therefore by reflexivity  $\langle \mu \rangle \mathcal{X} \sqsupset \mathcal{X}$ . Repeating this step we get  $\langle \mu \rangle \langle \mu \rangle \mathcal{X} \sqsupset \mathcal{X}$  that is  $\langle \mu^2 \rangle \mathcal{X} \sqsupset \mathcal{X}$ , etc.

We have  $\langle \mu^n \rangle \mathcal{X} \sqsupset \mathcal{X}$  and thus  $\langle \mu^n \rangle \mathcal{X} \neq \mathcal{Y}$  that is  $\mathcal{X} [\mu^n] \mathcal{Y}$ .  $\square$

EXAMPLE 2378. Connected funcoid without a path between given ultrafilters.

PROOF. Consider  $|\mathbb{R}|$ . It is connected (prove!) but there is no path (prove!) between two distinct singletons.  $\square$

THEOREM 2379. If meet of two connected (regarding a funcoid) filters is nonempty, then their join is connected.

PROOF. Let  $\mathcal{A}$  and  $\mathcal{B}$  be intersecting filters, both connected regarding an endofuncoid  $\mu$ . Let  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A} \sqcup \mathcal{B}$  for proper filters  $\mathcal{X}, \mathcal{Y}$ . Then either  $\mathcal{X}$  or  $\mathcal{Y}$  intersects  $\mathcal{A} \cap \mathcal{B}$ . Without loss of generality assume  $\mathcal{X} \cap \mathcal{A} \cap \mathcal{B} \neq \perp$ . Also  $\mathcal{Y}$  intersects either  $\mathcal{A}$  or  $\mathcal{B}$ . Without loss of generality assume  $\mathcal{Y} \cap \mathcal{A} \neq \perp$ .

Note  $\mathcal{X} \cap \mathcal{A} \neq \perp$ .

We have  $(\mathcal{X} \cap \mathcal{A}) \sqcup (\mathcal{Y} \cap \mathcal{A}) = (\mathcal{X} \sqcup \mathcal{Y}) \cap \mathcal{A} = (\mathcal{A} \sqcup \mathcal{B}) \cap \mathcal{A} = \mathcal{A}$ . So  $\mathcal{X} \cap \mathcal{A} [\mu] \mathcal{Y} \cap \mathcal{A}$  because  $\mathcal{A}$  is connected, consequently  $\mathcal{X} [\mu] \mathcal{Y}$  that is  $\mathcal{A} \sqcup \mathcal{B}$  is connected.  $\square$

THEOREM 2380. If meet of two connected (regarding a reloid) filters is nonempty, then their join is connected.

PROOF. Let  $S_1^*(\mu \cap (\mathcal{A} \times \mathcal{A})) = \mathcal{A} \times \mathcal{A}$ ;  $S_1^*(\mu \cap (\mathcal{B} \times \mathcal{B})) = \mathcal{B} \times \mathcal{B}$  for filters  $\mathcal{A} \neq \mathcal{B}$ .

$S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) = S_1^*(\mu \cap ((\mathcal{A} \times \mathcal{A}) \sqcup (\mathcal{B} \times \mathcal{B}) \sqcup (\mathcal{A} \times \mathcal{B}) \sqcup (\mathcal{B} \times \mathcal{A}))) \supseteq S_1^*(\mu \cap (\mathcal{A} \times \mathcal{A})) \sqcup S_1^*(\mu \cap (\mathcal{B} \times \mathcal{B})) \supseteq (\mathcal{A} \times \mathcal{A}) \sqcup (\mathcal{B} \times \mathcal{B})$ .

Let for example  $x \in \text{atoms } \mathcal{A}$ . Then  $\langle S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) \rangle x \supseteq \mathcal{A}$  and (taking into account  $\mathcal{A} \neq \mathcal{B}$ ):

$$\langle \mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B})) \rangle \langle S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) \rangle x \supseteq \mathcal{B}.$$

Thus  $\langle S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) \rangle x \supseteq \mathcal{A}$  and  $\langle S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) \rangle x \supseteq \mathcal{B}$  for every ultrafilter  $x \in \text{atoms}(\mathcal{A} \sqcup \mathcal{B})$ , that is  $\langle S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) \rangle x \supseteq \mathcal{A} \sqcup \mathcal{B}$ . So  $S_1^*(\mu \cap ((\mathcal{A} \sqcup \mathcal{B}) \times (\mathcal{A} \sqcup \mathcal{B}))) \supseteq \mathcal{A} \sqcup \mathcal{B}$  that is  $\mathcal{A} \sqcup \mathcal{B}$  is connected.  $\square$

COROLLARY 2381. Distinct connected components (for both a funcoid or a reloid) don't intersect.

PROOF. If connected components  $\mathcal{A} \neq \mathcal{B}$  intersect, then  $\mathcal{A} \sqcup \mathcal{B}$  is a connected filter and either  $\mathcal{A} \sqcup \mathcal{B} \sqsupset \mathcal{A}$  or  $\mathcal{A} \sqcup \mathcal{B} \sqsupset \mathcal{B}$  what contradicts to the definition of connected components.  $\square$

If we add the requirement  $\mathcal{X} \asymp \mathcal{Y}$  to the definition of connected regarding funcoid, it is nonequivalent. Proof??: Consider connectedness of an ultrafilter.

PROPOSITION 2382.  $S(\mu) = S_1(\mu \sqcup 1)$  if  $\mu$  is an endorelation, endofuncoid, or endoreloid. **FiXme:** for  $S^*$ , too.

PROOF. By proved above  $(\mu \sqcup 1)^n = 1 \sqcup \mu \sqcup \dots \sqcup \mu^n$ .

Thus  $S_1(\mu \sqcup 1) = (1 \sqcup \mu) \sqcup (1 \sqcup \mu \sqcup \mu^2) \sqcup \dots = 1 \sqcup \mu \sqcup \mu^2 \sqcup \dots = S(\mu)$ .  $\square$

**FiXme:** also algebraic properties of  $S_1$  and  $S_1^*$

THEOREM 2383. **FiXme:** Move this theorem in the book,  $\mathcal{X} [\prod S] \mathcal{Y} \Leftrightarrow \forall f \in S : \mathcal{X} [f] \mathcal{Y}$  if  $S$  is a generalized filter base.

PROOF.  $\mathcal{X} [\prod S] \mathcal{Y} \Leftrightarrow (\mathcal{X} \times^{\text{FCD}} \mathcal{Y}) \sqcap \prod S \neq \perp \Leftrightarrow \prod_{f \in S} f \sqcap (\mathcal{X} \times^{\text{FCD}} \mathcal{Y}) \neq \perp \Leftrightarrow$   
(by properties of generalized filter bases)  $\Leftrightarrow \forall f \in S : f \sqcap (\mathcal{X} \times^{\text{FCD}} \mathcal{Y}) \neq \perp \Leftrightarrow \forall f \in S : \mathcal{X} [f] \mathcal{Y}$ .  $\square$

THEOREM 2384. The following are pairwise equivalent for a funcoid  $\mu$  and filter  $\mathcal{A}$ :

- 1°.  $\mathcal{A}$  is connected regarding funcoid  $\mu$
- 2°.  $\mathcal{A}$  is connected regarding every funcoid in  $\text{up } \mu$ .
- 3°.  $\mathcal{A}$  is connected regarding every funcoid in  $\text{up}^\Gamma \mu$ .

PROOF. TODO: “Connectedness” should be moved after “Funcoids are filters” to use  $\Gamma$  in this proof.

1 $\Rightarrow$ 2 $\Rightarrow$ 3. Obvious.

3 $\Rightarrow$ 1. Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{F}(\text{Ob } \mu)$  and  $\mathcal{X} \sqcup \mathcal{Y} = \mathcal{A}$ . Then  $\forall f \in \text{up}^\Gamma \mu : \mathcal{X} [f] \mathcal{Y}$ . Therefore by the theorem ??  $\mathcal{X} [\prod \text{up}^\Gamma \mu] \mathcal{Y}$  that is  $\mathcal{X} [\mu] \mathcal{Y}$ . So  $\mathcal{A}$  is connected regarding  $\mu$ .  $\square$

CONJECTURE 2385. For a **Rel**-morphism  $F$  and a filter  $\mathcal{A}$  the following are pairwise equivalent:

- 1°.  $\mathcal{A}$  is connected regarding  $\uparrow^{\text{FCD}} F$ .
- 2°.  $\mathcal{A}$  is connected regarding  $\uparrow^{\text{RLD}} F$ .
- 3°. there is a  $F$ -path in  $\mathcal{A}$  for every two ultrafilters  $a, b \in \text{atoms } \mathcal{A}$ .

Proposed counterexample against  $\mathcal{A}$  is connected regarding  $f$  iff it is connected regarding  $(\text{FCD})f$ :  $f = \mathcal{A} \times_F^{\text{RLD}} \mathcal{A}$ . First calculate  $(\mathcal{B} \times_F^{\text{RLD}} \mathcal{C}) \circ (\mathcal{A} \times_F^{\text{RLD}} \mathcal{B})$  (and also for oblique product).

Trying to calculate  $(\mathcal{B} \times_F^{\text{RLD}} \mathcal{C}) \circ (\mathcal{A} \times_F^{\text{RLD}} \mathcal{B})$ :

LEMMA 2386. There are such filters  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  and binary relation  $h$  that

$$h \sqsupseteq \mathcal{A} \times^{\text{FCD}} \mathcal{C} \wedge \neg \exists g \in \mathbf{Rel} : (g \sqsupseteq \mathcal{B} \times^{\text{FCD}} \mathcal{C} \wedge h \sqsupseteq g \circ (\mathcal{A} \times^{\text{FCD}} \mathcal{B})).$$

PROOF. Take  $\mathcal{A}$  a principal filter,  $\mathcal{B}$  a trivial ultrafilter and  $h \sqsupseteq \mathcal{A} \times^{\text{FCD}} \mathcal{C}$  such that  $h \not\sqsubseteq \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{C})$ . (It exists because  $\mathcal{A} \times^{\text{RLD}} \mathcal{C} \neq \mathcal{A} \times_F^{\text{RLD}} \mathcal{C}$ .)

Suppose that  $g \sqsupseteq \mathcal{B} \times^{\text{FCD}} \mathcal{C}$ . Then there is  $C \in \text{up } \mathcal{C}$  such that  $g \sqsupseteq \mathcal{B} \times C$ . Therefore  $g \circ (\mathcal{A} \times^{\text{FCD}} \mathcal{B}) = \mathcal{A} \times^{\text{FCD}} \langle g \rangle \mathcal{B} \sqsupseteq \mathcal{A} \times^{\text{FCD}} C = \mathcal{A} \times C$ .

But  $h \not\sqsubseteq \text{up}(\mathcal{A} \times^{\text{RLD}} \mathcal{C}) = \text{up}(\mathcal{A} \times C)$ . Thus  $h \not\sqsupseteq g \circ (\mathcal{A} \times^{\text{FCD}} \mathcal{B})$ .  $\square$

COROLLARY 2387. There are such filters  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  and binary relation  $h$  that

$$h \sqsupseteq \mathcal{A} \times^{\text{FCD}} \mathcal{C} \wedge \neg \exists f, g \in \mathbf{Rel} : (f \sqsupseteq \mathcal{A} \times^{\text{FCD}} \mathcal{B} \wedge g \sqsupseteq \mathcal{B} \times^{\text{FCD}} \mathcal{C} \wedge h \sqsupseteq g \circ f).$$

PROPOSITION 2388.  $(\mathcal{B} \times_F^{\text{RLD}} \mathcal{C}) \circ (\mathcal{A} \times_F^{\text{RLD}} \mathcal{B}) \neq \mathcal{A} \times_F^{\text{RLD}} \mathcal{C}$  for some proper filters  $\mathcal{A}, \mathcal{B}, \mathcal{C}$ .

PROOF. **FiXme: The proof is erroneous.**

Take (lemma)  $h \in \text{up}(\mathcal{A} \times_F^{\text{FCD}} \mathcal{C})$  such that for every  $f \in \text{up}(\mathcal{A} \times_F^{\text{RLD}} \mathcal{C})$ ,  $g \in \text{up}(\mathcal{B} \times_F^{\text{RLD}} \mathcal{C})$  we have  $h \not\sqsupseteq g \circ f$ .

We have  $h \in \text{up}(\mathcal{A} \times_F^{\text{RLD}} \mathcal{C})$  and for every  $f \in \text{up}(\mathcal{A} \times_F^{\text{RLD}} \mathcal{C})$ ,  $g \in \text{up}(\mathcal{B} \times_F^{\text{RLD}} \mathcal{C})$  we have [error]  $h \not\sqsupseteq g \circ f$ .

Thus  $\text{up}((\mathcal{B} \times_F^{\text{RLD}} \mathcal{C}) \circ (\mathcal{A} \times_F^{\text{RLD}} \mathcal{B})) \neq \text{up}(\mathcal{A} \times_F^{\text{RLD}} \mathcal{C})$ .  $\square$

## Relationships are pointfree funcoids

**THEOREM 2389.**  $((\text{FCD}), (\text{RLD})_{\text{in}})$  are components of a complete pointfree funcoid.

**PROOF.** For every ultrafilters  $x$  and  $y$  we have  $x [( \text{FCD} )( f \sqcap (\text{RLD})_{\text{in}} g )] y \Leftrightarrow x \times^{\text{RLD}} y \not\neq f \sqcap (\text{RLD})_{\text{in}} g \Leftrightarrow x \times^{\text{RLD}} y \sqsubseteq (\text{RLD})_{\text{in}} g \wedge x \times^{\text{RLD}} y \not\neq f \sqcap (\text{RLD})_{\text{in}} g \Leftrightarrow x \times^{\text{FCD}} y \in \text{atoms } g : x \times^{\text{RLD}} y \not\neq f \sqcap (\text{RLD})_{\text{in}} g \Leftrightarrow x \times^{\text{FCD}} y \in \text{atoms } g : x \times^{\text{RLD}} y \not\neq f \Leftrightarrow x \times^{\text{FCD}} y \in \text{atoms } g \wedge x \times^{\text{FCD}} y \sqsubseteq (\text{FCD})f \Leftrightarrow x [g \sqcap (\text{FCD})f] y$ .

Thus  $(\text{FCD})(f \sqcap (\text{RLD})_{\text{in}} g) = g \sqcap (\text{FCD})f$ . Consequently  $f \sqcap (\text{RLD})_{\text{in}} g = \perp \Leftrightarrow g \sqcap (\text{FCD})f = \perp$  that is  $g \not\neq (\text{FCD})f \Leftrightarrow f \not\neq (\text{RLD})_{\text{in}} g$ .

It is complete by theorem 1095.  $\square$

We will also prove in another way that  $(\text{FCD}), (\text{RLD})_{\text{in}}$  are components of pointfree funcoids:

**THEOREM 2390.**  $(\text{RLD})_{\text{in}}$  is a component of a pointfree funcoid (between filters on boolean lattices).

**PROOF.** Consider the pointfree funcoid  $\mathcal{R}$  defined by the formula  $\langle \mathcal{R} \rangle^* F = (\text{RLD})_{\text{in}} F$  for binary relations  $F$  (obviously it does exist). Then  $\langle \mathcal{R} \rangle f = \langle \mathcal{R} \rangle \sqcap^{\text{FCD}} \text{up}^\Gamma f = \sqcap_{F \in \text{up}^\Gamma f}^{\text{RLD}} \langle \mathcal{R} \rangle^* F = \sqcap_{F \in \text{up}^\Gamma f}^{\text{RLD}} (\text{RLD})_{\text{in}} F = (\text{RLD})_{\text{in}} \sqcap_{F \in \text{up}^\Gamma f}^{\text{FCD}} F = (\text{RLD})_{\text{in}} f$ .  $\square$

**THEOREM 2391.**  $(\text{FCD})$  is a component of a complete pointfree funcoid (between filters on boolean lattices).

**PROOF.** Consider the pointfree funcoid  $\mathcal{Q}$  defined by the formula  $\langle \mathcal{Q} \rangle^* F = (\text{FCD})F$  for binary relations  $F$  (obviously it does exist). Then  $\langle \mathcal{Q} \rangle f = \langle \mathcal{Q} \rangle \sqcap^{\text{RLD}} \text{up } f =$  (because  $\text{up } f$  is a filter base)  $= \sqcap_{F \in \text{up } f}^{\text{FCD}} \langle \mathcal{Q} \rangle^* F = \sqcap_{F \in \text{up } f}^{\text{FCD}} (\text{FCD})F = \sqcap_{F \in \text{up } f}^{\text{FCD}} F = \sqcap^{\text{FCD}} \text{up } f = (\text{FCD})f$ .  $\square$

**PROPOSITION 2392.**  $(\text{FCD}) \sqcap S = \sqcap_{f \in S} (\text{FCD})f$  if  $S$  is a filter base of reloids (with the same sources and destinations).

**PROOF.** Theorem 839.  $\square$

**CONJECTURE 2393.**  $(\text{RLD})_{\text{in}} \sqcap S = \sqcap_{f \in S} (\text{RLD})_{\text{in}} f$  if  $S$  is a filter base of funcoids (with the same sources and destinations).

## Manifolds and surfaces

### 1. Sides of a surface

DEFINITION 2394. Let  $\mu$  be an endofunctor on a set  $U$ . *Surface side* of a set  $T \subseteq \text{Ob } \mu$  is a connected component (regarding  $\mu$ ) of the filter  $(\langle \mu \rangle^* T) \setminus T$ . **FiXme:**  $\mu$  is used twice in this definition. We may generalize for two different functors instead.

Keep in mind that the above definition may work nicely if  $\mu$  is a complete functor induced by a topological space.

EXAMPLE 2395. For an  $\mathbb{R}^{n-1}$  subspace  $T$  of a  $\mathbb{R}^n$  ( $n \geq 1$ ) euclidean space and the complete functor  $\mu$  induced by the usual topology:

- 1°.  $T$  has exactly two surface sides.
- 2°. The filter  $\langle \mu \rangle^* \{a\} \setminus T$  (for every  $a \in T$ ) has exactly two connected components.

PROOF. Without loss of generality assume that

$$T = \left\{ \frac{(x_0, x_1, \dots, x_{n-2}, 0)}{x_0, x_1, \dots, x_{n-2} \in \mathbb{R}} \right\}; \quad a = (0, \dots, 0).$$

We have

$$\langle \mu \rangle^* \{a\} = \left( \uparrow \left\{ \frac{v \in \mathbb{R}^n}{v_{n-1} > 0} \right\} \cap \langle \mu \rangle^* \{a\} \right) \sqcup \left( \uparrow \left\{ \frac{v \in \mathbb{R}^n}{v_{n-1} < 0} \right\} \cap \langle \mu \rangle^* \{a\} \right).$$

Let us prove that  $\uparrow \left\{ \frac{v \in \mathbb{R}^n}{v_{n-1} > 0} \right\} \cap \langle \mu \rangle^* \{a\}$  and  $\uparrow \left\{ \frac{v \in \mathbb{R}^n}{v_{n-1} < 0} \right\} \cap \langle \mu \rangle^* \{a\}$  are connected components.

??

□

**1.1. Special points.** We will start from the example of open  $T = \left\{ \frac{(x, y, 0)}{x^2 + y^2 < 1} \right\}$  and closed  $T = \left\{ \frac{(x, y, 0)}{x^2 + y^2 \leq 1} \right\}$  disks in  $\mathbb{R}^3$ .

EXERCISE 2396. Prove that open disk (in a usual 3-dimensional space) has two surface sides and closed disk has one surface side.

### 2. Special points

DEFINITION 2397. *Surface cardinality* of a point  $a$  (an element of the set  $\text{Ob } \mu$ ) is the cardinality of the set of connected components of the filter  $\langle \mu \rangle^* \{a\} \setminus T$ .

DEFINITION 2398. *Cardinality regular point* is a point  $a$ , which has a neighborhood ( $X \in \text{up } \langle \mu \rangle^* \{a\}$ ) such that all points  $x \in X \cap T$  are of the same surface cardinality as the point  $a$ .

*Cardinality special point* is a point which is not cardinality regular.

DEFINITION 2399. *Isomorphism regular point* is a point  $a$ , which has a neighborhood ( $X \in \text{up } \langle \mu \rangle^* \{a\}$ ) such that for all points  $x \in X \cap T$  the filter  $\langle \mu \rangle^* \{a\}$  is isomorphic to  $\langle \mu \rangle^* \{x\}$ .

*Isomorphism special point* is a point which is not isomorphism regular.




FIGURE 15. Examples of surface cardinality

**FixMe:** Try to replace isomorphism  $f$  with some kind of filter embedding.

Consider the dihedral angle  $T$  produced by two half-planes. Are the points of intersection of the half-planes isomorphism-special? (They should not be considered special. If they are special, this is a probably flaw in the definition of isomorphism special.)

Consider union  $T$  of two intersecting lines on a plane. The intersection may be considered as a special point, because it has more connected components than the rest. We don't want to consider it special, however. We can restrict to consider special only points which have less connected components (rather than more) to correct this trouble. Also try to define it with some kind of morphisms of filters instead of isomorphism as in isomorphism-special.

EXERCISE 2400. Excluding special points (either cardinality or isomorphism) from closed disk produces open disk.

Let us note that special points of closed disk have surface cardinality 1 which is less than surface cardinality (2) of regular points. So, it is a conceivable idea to consider special points which have lesser surface cardinality than nearby points.

Consider the following two subsets of a plane (the lines are the set  $T$ , the small black blob is the point  $a$ , and the cyan blob symbolizes the filter  $\langle \mu \rangle^* \{a\} \setminus T$ ):

For one of the sets surface cardinality of  $a$  is 3 and for another it is 2.

Now define *shift special points*.

Let  $I$  be an interval on  $\mathbb{R}$  (containing zero?)

A point  $a$  is *shift special* if there exists a transformation (that is a continuous function  $f : I \times \mu \rightarrow \mu$  such that:

- 1°.  $f(0)$  is identity. **FixMe:** Is this condition needed?
- 2°. for every sufficiently small  $\epsilon > 0$  we have  $f(\epsilon, a) \in T$ ;
- 3°. there is  $\epsilon > 0$  such that for every  $0 < \epsilon' < \epsilon$  we have  $f(\epsilon')$  being not continuous at  $a$  regarding complete funcooid defined by the function  $x \mapsto \langle \mu \rangle^* \{x\} \setminus T$ .

We may consider to additionally require that every  $f(\epsilon)$  is isomorphism of funcooids.

EXAMPLE 2401.  $T$  is disk  $\left\{ \frac{(x,y,0)}{x^2+y^2 \leq 1} \right\}$ .  $f$  is the contraction  $(\epsilon, v) \mapsto \frac{1}{1+\epsilon}v$ .  $a = (1, 0, 0)$ .

In the usual topology  $f$  is continuous. In  $x \mapsto \langle \mu \rangle^* \{x\} \setminus T$  we have the function  $\epsilon \mapsto f(\epsilon)$  not continuous at zero. So  $a$  is a shift special point.

PROOF.  $f(0)(v) = v$ . Thus  $\langle f(0) \rangle (\langle \mu \rangle^* \{a\} \setminus T) = \langle \mu \rangle^* \{a\} \setminus T$  intersects the plane  $Z = 0$ . But  $f(0, a)$

?? □

QUESTION 2402. Can we exclude real numbers from the play?

QUESTION 2403. How cardinality special points, isomorphism special points and shift special points are related with each others?



QUESTION 2404. How the number of surface sides is related with usual surface sides for manifolds? [https://en.wikipedia.org/wiki/Orientability#Orientability\\_of\\_manifolds](https://en.wikipedia.org/wiki/Orientability#Orientability_of_manifolds)

REMARK 2405. Manifolds have no special points. (Prove!)

Prove that 2-manifold image which special points removed has the same number of sides as the defined above.

Another way to define special points: A special point is a point such that  $T\pi\langle\mu\rangle\{a\}$  is not isomorphic to  $T\pi\langle\mu\rangle\{x\}$  for nearby points  $x$ . Consider replacement of isomorphism with injection, surjection, etc. here and above.

How many sides has in  $\mathbb{R}^3$  a plane without one point?

Easy way to spot special points: They are boundary points in the topology (or funoid) induced on  $T$ . Alternatively we can consider points whose neighborhood in  $T$  is different (as non-isomorphic or maybe non-injective or non-surjective or like this) than of nearby points. Thus another way to remove special points: use interior funoid.

<https://math.stackexchange.com/q/2836833/4876>

## Bibliography

- [1] T. Banach and A. Ravsky. Each regular paratopological group is completely regular. *ArXiv e-prints*, October 2014.
- [2] Victor Porton. *Algebraic General Topology. Volume 1*. 2014.
- [3] Zahava Shmuely. The tensor product of distributive lattices. *algebra universalis*, 9(1):281–296.
- [4] Zahava Shmuely. The structure of galois connections. *Pacific J. Math.*, 54(2):209–225, 1974.